

To Study and Design of Photocatalytic Reactor for Removal of Organic Effluents from Wastewater

Aher Prasad Shrikrushna

Pravara Rural Education Society's, Sir Visvesvaraya Institute of Technology, Department of Chemical Engineering Nashik-422101

Abstract: *Few years ago, growing human needs led to industrialization providing amenities to make human life straightforward. However, due to the industrial growth, the pollution of air and water also reached to a degree where new and innovative processes/technologies are required to be developed. Therefore, varieties of techniques have been developed for pollution abatement. Water pollution from emerging contaminants (ECs) or emerging pollutants is an important environmental problem. Heterogeneous photocatalytic treatment, as advanced oxidation treatment of wastewater effluents, has been proposed to solve this problem. Heterogeneous photocatalytic process was studied for emergent contaminants removal using paracetamol as a model contaminant molecule. The oxidation of organic substrates represents one of the most important industrial chemical reactions, explaining the significant efforts invested in the research and development of new heterogeneous catalysts with increased activities and selectivities in these type reactions.*

Keywords: photo catalysis, oxidation, effluent, wastewater, pollutant.

1. Introduction

Current journalism has verified on a laboratory scale the probable of this shows potential technology to completely demolish organic pollutants dissolved or dispersed in water into harmless substances. While the devastation rate of *E. coli* was exaggerated by pH, dissolve oxygen (DO) and temperature (T), it was found that number of bacteria shattered throughout disinfection mainly depends on occurrence light intensity (I₀) and irradiation time (t), which follows an exponential connection. In semiconductor photocatalysis, the light absorbing species is a semiconducting material. The electronic structure of most semiconductor materials comprise a highest occupied band full of electrons called the valance band (VB), and a lowest untenanted band called the conductance band (CB). These bands are alienated by a Region that is principally devoid of energy levels and the disparity in energy between the two bands is called the band gap energy, E_{bg}. present at the surface, i.e. A, then the photogene rated conductance band electrons can respond with it to produce a reduced product, A⁻. The overall reaction can be summarized as follows:



In the sanitization of water via semiconductor photocatalysis, the electron acceptor, A, is habitually dissolved oxygen, and the electron donor, D, is the pollutant, which is typically unrefined. Under these situation the overall process is the semiconductor photocatalysed oxidative mineralisation of the natural pollutant by dissolved oxygen, and can be represent by the following equation: Pollutant O semiconductor minerals



where „minerals“ are CO₂, H₂O and, where appropriate, inorganic acids or salts, such as HCl or NaCl.

2. Literature Survey

2.1 Ashok Kumar

Intentional an general idea on semiconductor particulate system for hydrogen creation. Potentials of various photocatalysts for hydrogen production, such as CdS, TiO₂ and SiC, were discussed. furthermore unlike catalyst amendment methods, such as noble metal loading, toting up of sacrificial reagent and dye were evaluate given that then, a number of innovative technology have been urbanized excluding anion doping and metal ion-scion. This paper aims to give a wide-ranging review on the modern maturity of photocatalytic hydrogen invention using photocatalyst TiO₂. compare with other photocatalysts, TiO₂ is much more hopeful as it is steady, non-corrosive, environmentally friendly, plentiful and cost useful. Based on recent research works, the augmentation methods for photocatalytic hydrogen production are reviewed and summarized.

2.2 Gianluca Li Puma, Po Lock Yue

Intentional the application of photo catalysis for water treatment and purification on an industrial scale can be accelerate by the progress of both new photo reactor design and arithmetical models. A work of fiction, pilot-plant, thin-film, slurry photo catalytic reactor for water Treatment and purification is obtainable in this paper. The reactor is a 'spout photo catalytic reactor, consisting of a grace with your presence water bell irradiated from above. Such a reactor arrangement is particularly appropriate for large-scale solar applications of photo catalysis.

A dimensionless arithmetical model for the cascade photo catalytic reactor is presented in this paper. The model was urbanized using parameter that can be predictable effortlessly from genuine systems and replica solutions can be obtain with modest computational endeavor. The model was legalize with untried results from the photo catalytic oxidation of salicylic acid in a pilot-scale reactor using titanium dioxide (TiO₂) as the photo catalyst experiment

were perform under a number of deferent circumstances by anecdotal substrate meditation, concentration of the occasion radiation, catalyst loading, flow rate, salvage ratio and water spray diameter. The model results were found to healthy the new data well in all cases. The model draw near can be extensive to other thin-film slurry photo catalytic reactors.

2.3 William A. Jacoby

Studied article will review the work that has been available on disinfection and the butchery of cancer cells using photocatalytic chemistry with titanium dioxide (TiO₂). This is an application of photocatalytic chemistry that has been under vigorous examination since 1985. Because the nature of the research is such that it brings together unrelated discipline this review provides background on photocatalytic chemistry, essential characteristics of intention organisms, potential applications, and the toxicology of titanium dioxide

3. Types of photoreactor

3.1 Optical Fiber Tube Reactor

Impulsive organic compounds (VOCs) are distinctive gaseous emission from a range of industries and pretense hazard for human health and environment a lot of technologies presently used for the treatment of VOCs, such as adsorption and scrubbing processes, merely migrate the pollutants from gas phase to another phase. Oxidation, such as ultraviolet/titanium process, is measured to be an efficient process for decomposing volatile organic pollutants in gaseous phase.

In recent years, many research have studied the likelihood and reaction mechanism of TiO₂-based photo catalytic breakdown of several gas-phase organic compounds employ fixed-bed photo reactors operated in either batch or permanent flow modes..




Figure: Typical Type Photocatalytic Reactor

3.2 Packed bed Reactors

In this type of reactors the fluid stream to be treated flows during the packed bed, comes into speak to with the irradiated particles holding the TiO₂. Some of the drawbacks that this design may tolerate are a low surface area to reactor volume ratio and low use of irradiation and this in view of both light absorption and scattering. Different types of fixed bed reactors have been projected: flat or curved walls, grooved walls, monoliths, packed beds and reticulated configurations. Changrani and Raupp proposed a reticulated foam photocatalytic reactor and solved the representation with Monte Carlo simulation; these authors also used a deterministic two-dimensional (2-D) assorted model to simulate this reactor. At the same time, a model for absorption by TiO₂ films in a corrugated-plate photocatalytic reactor was described by Zhang etc.




Figure: Packed Bed Photocatalytic reactors

3.3 Spinning Disc Reactor

The rotor-stator spinning disc reactor consists of a rotating disc in a cylindrical lodging. The distance between the rotor and the reactor wall, the stator, is small, in the order of 1 mm. Two reactor configurations are studied. In the first configuration, the liquid, which is the continuous phase, is injected to the reactor from the top. Gas bubbles are sheared off at the gas inlet, due to the high velocity gradient, and thus shear force, between the rotor and the stator. The gas

bubble size reduces with increasing rotational disc speed.




Figure: Spinning Disc Reactor Photocatalysis mechanism




Figure: Mechanism of photocatalyst

4.2 Mechanism

When photocatalyst titanium dioxide (TiO_2) absorbs Ultraviolet (UV) radiation from sunlight or illuminated light preliminary place (fluorescent lamps), it will engender pairs of electrons and holes. The electron of the valence band of titanium dioxide becomes thrilled when illuminate by light. The excess energy of this active electron promoted the electron to the conveyance band of titanium dioxide therefore creating the negative-electron (e^-) and positive-hole (h^+) pair. This phase is referred as the semiconductor's 'photo-excitation' state. The energy difference between the valence band and the conduction band is known as the 'Band Gap'. Wavelength of the light essential for photo-excitation is: $1240 \text{ (Planck's constant, } h) / 3.2 \text{ eV (band gap energy) = } 388 \text{ nm}$. **7.4 Super-Hydrophilic**

When the exterior of photocatalytic film is uncovered to light, the get in touch with angle of the photocatalyst surface with water is abridged steadily. After enough introduction to light, the surface reaches super-hydrophilic. In other words, it does not ward off water at all, so water cannot survive in the shape of a drop, but spreads utterly on the surface of the substrate. And the water took the form of a highly standardized thin film, which behaves optically like a apparent sheet of glass. The hydrophilic nature of titanium dioxide, fixed with the gravity, will permit the dust particles to be swept away following the water stream, thus making the product self-cleaning.

The positive-hole of titanium dioxide breaks at a distance

4. Definition of Photocatalyst

4.1 Photo-Catalysis is termed as "acceleration by the presence of as catalyst". A catalyst does not change in itself or being obsessive in the chemical reaction. This definition includes photosensitization, a process by which a photochemical amendment occurs in one molecular entity as a result of preliminary absorption of radiation by another molecular entity called the photosensitized. Chlorophyll of plants is a type of photocatalyst. Photocatalysis compared to photosynthesis, in which chlorophyll captures sunlight to turn water and carbon dioxide into oxygen and glucose, photocatalysis form strong oxidation agent to breakdown any organic matter to carbon dioxide and water in the presence of photocatalyst, light and water.

the water molecule to form hydrogen gas and hydroxyl radical. The negative-electron reacts with oxygen molecule to form super oxide anion. This cycle continues when light is obtainable. The overall mechanism of photocatalytic reaction of titanium dioxide.

4.3 Photocatalytic Oxidation

The most authoritative advanced oxidation systems are based on the generation of hydroxyl radicals. The hydroxyl radical is an tremendously powerful oxidation agent, second only to Fluorine in power (2.23 in Relative Oxidizing Power). Following is a listing of common chemical oxidants, placed in the order of their oxidizing strength:

4.4 Deodorizing Effect

On the deodorizing application, the hydroxyl radicals accelerate the breakdown of any Volatile Organic Compounds or VOCs by destroying the molecular bonds. This will help merge the organic gases to form a single molecule that is not harmful to humans thus enhance the air cleaning efficiency. Some of the examples of odor molecules are: Tobacco odor, formaldehyde, nitrogen dioxide, urine and fecal odor, gasoline, and many other hydro carbon molecules in the atmosphere. Air purifier with TiO_2 can prevent smoke and soil, pollen, bacteria, virus and harmful gas as well as seize the free bacteria in the air by filtering percentage of 99.9% with the help of the highly oxidizing effect of photocatalyst (TiO_2).

4.5 Anti-Bacterial Effect

Photocatalyst does not only slay bacteria cells, but also decay the cell itself. The titanium dioxide photocatalyst has been found to be more effective than any other antiseptic agent, because the photocatalytic reaction works even when there are cells covering the surface.

5. Future Outlook

There is a enormous body of literature, including a number of appraisals, recitation TiO₂ catalysed photochemistry [18-20]. Noteworthy are the 1995 contribution of Hoffmann et al. which includes 441 references and the excellent review paper by Hashimoto et al. [15] This study confirms that photocatalysis is fitting not only for decolorizing colored aqueous effluents but also promisingly degrading the dyes and its poverty intermediates to harmless mineral products (CO₃, H₃O, SO₄'etc). With good photocatalytic activity under UV irradiation and the ability to be readily separated from the reaction system, this novel kind of catalyst exhibited the probable to be effective in the action of organic pollutants in aqueous systems. ia are actively propagating. The end toxin produced at the death of cell is also expected to be decomposed by photocatalytic action. Titanium dioxide does not deteriorate and it shows a long-term anti-bacterial effect. Generally speaking, disinfections by titanium oxide is three times stronger than chlorine, and 1.5 times stronger than ozone.

6. Result

Focuses on the collection of the best route by which TiO₂ catalysts were prepared, based on the application of the catalyst on the degradation of Methylene Blue (MB), a common unrefined dye, by varying all possible parameters. It is inferred that the photochemical degradation of methylene blue occurred effectively over the prepared catalysts. The percentage degradation of the dye for various systems could be correlated with the large surface area, crystallite size, mesoporous nature and the band gap.

7. Acknowledgement

I take great pleasure in presenting this paper on “**To study and design of photocatalytic reactor for removal of organic effluents from wastewater**” Foremost, I would like to express my sincere gratitude to all the staff members of CHEMICAL DEPARTMENT for the support and help which they always provided us with whenever it was required. I would like to thanks my colleagues who work with me through thick and thin for successful completion of paper. Last but not the list; I would like to thanks my mother & father they always providing support & ,encouraging me.

References

- [1] C.S.turchi and D.w ollis, J. catalysis 122(1990),178.
- [2] Studies on catalysis by titania Joyes Jacob.
- [3] C.A martin, G.Camera-roda and f.sntarelli, catalysis Today 48(1999),307.

- [4] A.Alexiadis, G Daldi and mazzarino catalysis Today 66(2001),467
- [5] H.y.chen, o.zahraa, Douchdy, f thomos and J.y.Bottero (1995),1279.
- [6] D.chem, F li and A.k Ray catalysis today 66(2001),475
- [7] Inhidoct,sco-keyn leeand Andrew mills.
- [8] D.S.ollis and turchi Environ.prog,229(1990)
- [9] D.S.Bhthande, V.G Pangarkar and A.A. Beenackers, J.chem technol.Biotechnol 77,162(2000)
- [10] J.galvezands.N Rodriguez online publication:- <http://www.unesco.org>(2002)
- [11] G.Li Puma and p.Lyue, chem. Engg.sci.53(1998),2993
- [12] E.piera,M.L.Tejedor,M.E.Zorn,M.A.Andorson,Appl. Cotal,B 46(2003)671.
- [13] A.L.Linebigler, G.L,Jtyater Jr.chem Rev.95(1995)735-58
- [14] K. Hashimoto, H.Irie, fusishima A.Jpn J Appl.phys part 1.44 (2005)8269.85.
- [15] Hoffmann et al.1995