

(σ, τ) –Lie Ideals with Some New Circulars

Majid Neamah^{1*}, Hala Mehdy²

¹University of Baghdad, College of Science, Mathematics Department, Baghdad, Iraq

²University of Mustansiriyah, Baghdad, Iraq

¹mmathemtic[at]yahoo.com, ²hmprogram[at]yahoo.com

Abstract: Let R be a ring optional, U be an additive subgroup of R and $\sigma, \tau: R \rightarrow R$ be two mapping $[x, y]_{\sigma, \tau} = x\sigma(x) - \tau(y)x$ in most of our study we will consider R is prime ring with characteristic not equal 2, and (σ, τ) functions equivalent automorphsem. The basic aim of this study Is the study of circulating Lieidealto (σ, τ) - Lie ideal mainstreaming some results at (σ, τ) - Lie ideal using theorems non-generalized for the purpose of circulating provable and the use of these proofs generalized to help prove theorems other non-generalized. We can prove a lot of non-generalized theorems on the subject Lie ideal by these theorems that have been circulated in this research. We have used in this research

$$\{R \ni x \forall R; c(y) + \tau(x)c\} = C_{\sigma, \tau}$$

$$d(xy) = d(x)\sigma(y) + \tau(x)d(y), \forall x, y \in R$$

Results are as follows

1) **Theorem (2.1).** Let $d_1: R \rightarrow R$ be a (σ, τ) - derivation and $d_2: R \rightarrow R$ be an (α_j, α_j) - derivation and $d_3: R \rightarrow R$ be an (β, β) - derivation. Such that $d_3\beta = \beta d_3, d_2\beta = \beta d_2, d_1\beta = \beta d_1$, where β is automorphsim of R . If $U \neq (0)$ is an ideal of R such that $d_3(U) \subset U$ and $d_1d_2d_3(U) = 0$, Then either $d_1 = 0$ or $d_2 = 0$ or $d_3 = 0$.

2) **Theorem (2.2)**(In general). Let $d_1: R \rightarrow R$ be (σ, τ) - derivation and $d_i: R \rightarrow R$ be an (α_j, α_j) - derivation such that $d_1\alpha_1 = \alpha_1d_1, d_i\alpha_j = \alpha_jd_i, i = 2, \dots, n, j = n - 1$, Where is α_j an outomorphism of R if $U \neq (0)$ s an ideal of R such that $d_1(U) \subset U$ A $d_1d_2 \dots d_n(U) = (0), n \in \mathbb{N}$,

Then either $d_1 = 0$ or $d_2 = 0$ or $d_n = 0$

3) **Theorem (2.3).** Let U be nonzero ideal of R and $a, b, c \in U$,

if $[c, [a, [b, x]]]_{\sigma, \tau} = 0, \forall x \in U$ for all $x \in U$, then either

1. $c, a \in C_{\sigma, \tau}$ or $b \in Z(R)$.

2. $c \in C_{\sigma, \tau}$ or $a, b \in Z(R)$.

4) **Theorem (2.4)** (In general). Let U be nonzero ideal of R and,

if $a_1, a_2, \dots, a_n \in U, \forall x \in U$ then either

1. $a_n, a_{n-1}, \dots, a_2 \in C_{\sigma, \tau}$ or $a_1 \in Z(R)$

2. $a_n \in C_{\sigma, \tau}$ or $a_{n-1}, \dots, a_1 \in Z(R)$

Keywords: Lie ideals, ring, semi-ring, characteristic ring, derivation, homomorphic.

1. Introduction

This work is a continuation of a series results that have been obtained by some researchers (K. A. Jassim) and Dr. (A.A. Hameed) see [25-26].

Let R be a ring, U be an additi ve subgroup of R , U is called a lie ideal of R if $[U, R] \subset U$.

We generalized this definition that : If we have $\sigma, \tau: R \rightarrow R$ be two mapping then

1) U is called a (σ, τ) right Lie ideal of R if $[U, R]_{\sigma, \tau} \subset U$.

2) U is called a (σ, τ) lift Lie ideal of R if $[R, U]_{\sigma, \tau} \subset U$.

3) U is called a (σ, τ) -Lie ideal of R if U is both (σ, τ) -right Lie ideal and (σ, τ) -left Lie ideal of R .

This chapter consists of two sections, in section one we give the basic definitions and study the relations among them, we illustrate some of these results by some examples. In most of our study we will consider R is prime ring with characteristic not equal 2, and σ, τ functions equivalent automorphisms. In section two, we gave important results when U is a (σ, τ) right Lie ideal of R , such that if $[U, U]_{\sigma, \tau} \subset C_{\sigma, \tau}$ then either $U \subset Z(R)$ or $U \subset C_{\sigma, \tau}$. Also if U is a subring of R with new theorems which represents the generalization to lemma (3.1) when $n=3$ and when n in general, as well as we concluded a new theorem (3.3) we got some relations between the center of the R , denoted by $Z(R)$ and between the centralizer of x in R , denoted by $C_R(x)$.

The aim of the study

1) Study is a generalization of the concept is perfect for Lie ideal to (σ, τ) -Lie ideal and a generalization some of the results of the ideal for ideal to (σ, τ) -Lie ideal as well as provable.

2) To study the relationship between the derivative and the (σ, τ) -Lie ideal and give some important results, we have used in this study

$\{R \ni x \forall R; c\sigma(y) + \tau(x)c; R \in c\} = C_{\sigma, \tau}$ as well as we used the following formula to complete derivation process $d(xy) = d(x)\sigma(y) + \tau(x)d(y), \forall x, y \in R$.

2. Basic Concepts

Definition (2.1). A ring R is called a prime ring if $aRb = (0), a, b \in R$, implies that $a = 0$ or $b = 0$.

Definition (2.2). A ring R is called a semi prime ring if $aRb = (0), a, b \in R$, implies that $a = 0$.

Definition (3.3). Let R be an arbitrary ring. If there exists a positive integer n such that $na = 0$ for all $a \in R$, then the smallest positive integer with this property is called the characteristic of the ring, by symbols we write $chR = n$. If no such positive integer exists (that is, $n=0$ is the only integer for which $na = 0$, for all a in R), then R is said to be of characteristic zero.

Remark (2.1). We can show easily that if R is a prime ring with characteristic not equal n is equivalent to n -torsion free.

Definition (2.4). Let R be a ring. Define a Lie product $[,]$ on R as follows

$$[x, y] = xy - yx \forall x, y \in R.$$

Remark (2.2). Let R be a ring, then $\forall x, y \in R$ we have: -

$$[x, yz] = y[x, z] + [x, y]z$$

$$[x + y, z] = [x, z] + [y, z]$$

$$[xy, z] = x[y, z] + [x, z]y$$

Definition (2.5). Let A be a Lie subring of a ring R . An additive subgroup $U \subset A$ is said to be a Lie ideal of A , if whenever $u \in U$ and $a \in A$, then $[u, a] \in U$.

Definition (2.6). Let R be a ring Define the product $[x, y]_{\sigma, \tau} = x\sigma(y) - \tau(y)x, \forall x, y \in R$.

Remark (2.3). Let R be a ring and let $\sigma, \tau : R \rightarrow R$ be two mappings. The $\forall x, y, z \in R$, we have:

$$1) [x + y, z]_{\sigma, \tau} = [x, z]_{\sigma, \tau} + [y, z]_{\sigma, \tau}.$$

$$2) [xy, z]_{\sigma, \tau} = x[y, z]_{\sigma, \tau} + [x, (z)]_{\sigma, \tau} y = x[y, \sigma(z)] + [x, z]_{\sigma, \tau} y.$$

Remark (2.4). Let R be a ring and let $\sigma, \tau : R \rightarrow R$ be two homomorphism's. Then $\forall x, y \in R$, we have:

$$[x, yz]_{\sigma, \tau} = \tau(y)[x, z]_{\sigma, \tau} + [x, y]_{\sigma, \tau} \sigma(z)$$

Definition (2.7). Let R be a ring, U be an additive subgroup of R and, $\sigma, \tau : R \rightarrow R$ be two mappings. Then

- 1) U is called a (σ, τ) -right Lie ideal of R if $[U, R]_{\sigma, \tau} \subset U$.
- 2) U is called a (σ, τ) -left Lie ideal of R if $[R, U]_{\sigma, \tau} \subset U$.
- 3) U is called a (σ, τ) -Lie ideal of R if U is both (σ, τ) -right Lie ideal and (σ, τ) -left Lie ideal of R .

Definition (2.8). Let R be a ring, the center of R , denoted by $Z(R)$, is the set

$$\{a \in R; ar = ra \forall r \in R\}.$$

Definition (2.9). Let X be a nonempty subset of R , the centralizer of X in R , denoted, by $C_R(X)$, is the set.

$$\{a \in R; [x, a] = 0 \forall x \in X\}$$

Definition (2.10). Let R be a ring and let, $\sigma, \tau : R \rightarrow R$ be two mappings. (σ, τ) -centralizer of R , denoted by $C_{\sigma, \tau}$ is the set

$$\{c \in R; c\sigma(x) = \tau(x)c \forall x \in R\}.$$

Definition (2.11). Let R be a ring. An additive mapping $d: R \rightarrow R$ is called a derivation on R if

$d(xy) = d(x)y + xd(y) \forall x, y \in R$. We say that d is an inner derivation if there exists an element $a \in R$ such that

$$d(x) = [a, x] \forall x \in R.$$

Definition (2.12). Let R be a ring. An additive mapping $d : R \rightarrow R$ is called a (σ, τ) -derivation where, $\sigma, \tau : R \rightarrow R$ be two mappings, if $d(xy) = d(x)\sigma(y) + \tau(x)d(y) \forall x, y \in R$. It is clear that every derivation is a (σ, τ) -derivation.

Definition (2.13). Let $d : R \rightarrow R$ be an additive mapping then we say that d is a (σ, τ) -inner derivation if there exists an element $a \in R$ such that $d(x) = [a, x]_{\sigma, \tau}, \forall x \in R$

3. (σ, τ) -Right Lie ideals

The following lemmas help us to prove the main theorems

Lemma (3.1). Let d be a (σ, τ) -derivation of R and $a \in R, U$ be a nonzero ideal of R . If $(U) = (0)$ or $(d(U)a) = (0)$, then either $a = 0$ or $d = 0$.

Lemma (3.2). Let $d_1 = R \rightarrow R$ be a (σ, τ) -derivation and $d_2 = R \rightarrow R$ be an (α, α) derivation such that $d_2\alpha = \alpha d_2, d_1\alpha = \alpha d_1$, where α is an automorphism of R . If $U \neq (0)$ is an ideal of R such that $d_2(U) \subset U$ and $d_1d_2(U) = (0)$, then either $d_1 = 0$ or $d_2 = 0$.

Proof. For any $u, v \in U, uv \in U$. By hypothesis $d_1d_2(U) = (0)$, So,

$$\begin{aligned} 0 &= d_1d_2(uv) = d_1(d_2(u)\alpha(v) + \alpha(u)d_2(v)) = d_1(d_2(u)\alpha(v)) + d_1(\alpha(u)d_2(v)) \\ &= d_1(d_2(u))\sigma(\alpha(v)) + \tau(d_2(u))d_1(\alpha(v)) + d_1(\alpha(u))\alpha(d_2(v)) + \alpha(\alpha(u))d_1(d_2(v)). \end{aligned}$$

Since $d_1d_2(U) = (0)$, also $-\tau(d_2(u))d_1(\alpha(v)) + d_1(\alpha(u))\sigma(d_2(v)) = 0$, that is,

$$d_1(\alpha(u)\sigma(d_2(v)) + \tau(d_2(u))d_1(\alpha(v))) = 0, \forall u, v \in U \dots \dots \dots (1)$$

Replacing u by $d_2(u)$ in (1). We get

$$\begin{aligned} d_1(\alpha(d_2(u)))\sigma(d_2(v)) + \tau(d_2(d_2(u)))d_1(\alpha(v)) &= 0 \text{ Using } \alpha d_1 = d_1\alpha \text{ we have} \\ \tau(d_2^2(u))d_1(\alpha(v)) = 0, \forall u, v \in U \dots \dots \dots (2) \end{aligned}$$

Since α is an automorphism, hence α^{-1} exists such that by Lemma (2.1) we get

$$d_2^2(u) = 0 \forall u \in U \text{ or } d_1 = 0. \text{ That is } d_2^2(U) = (0) \text{ or } d_1 = 0.$$

Suppose $d_1 \neq 0$, then $d_2^2(U) = (0)$. For any $u, v \in U$, so $uv \in U$. Hence,

$$\begin{aligned} 0 &= d_2^2(uv) = d_2(d_2(uv)) = d_2(d_2(u)\alpha(v) + \alpha(u)d_2(v)) = d_2(d_2(u)\alpha(v)) + d_2(\alpha(u)d_2(v)) \\ &= d_2(d_2(u))\alpha(\alpha(v)) + \alpha(d_2(u))d_2(\alpha(v)) + d_2(\alpha(u))\alpha(d_2(v)) + \alpha(\alpha(u))d_2(d_2(v)). \end{aligned}$$

$$\text{But } d_2^2(U) = 0, \text{ we get } \alpha(d_2(u))d_2(\alpha(v)) + d_2(\alpha(u))\alpha(d_2(v)) = 0.$$

Also, $\alpha d_2 = d_2 \alpha$ we have $d_2(\alpha(u))d_2(\alpha(v)) + d_2(\alpha(u))d_2(\alpha(v)) = 0$. So $2d_2(\alpha(u))d_2(\alpha(v)) = 0$. Since R is a prime ring with $ChR \neq 2$, then $d_2(\alpha(u))d_2(\alpha(v)) = 0$. So, $\alpha(d_2(u)d_2(v)) = 0$ and $d_2(u)d_2(v) = 0, \forall u, v \in U$. Therefore $d_2(U)d_2(U) = (0)$.

By Lemma (2.1), we get either $d_2(U) = (0)$ or $d_2 = 0$.

If $d_2(U) = (0)$, then $d_2(ru) = 0, u \in U, r \in R$. This implies

$0 = d_2(r)\alpha(ru) = \alpha(r)d_2(u) = d_2(r)\alpha(u)$. That is, $d_2(r)\alpha(u) = 0 \forall u \in U, r \in R$. Now

$0 = d_2(r)\alpha(ru) = d_2(r)\alpha(r)\alpha(u) \forall u \in U, r \in R$.

Since R is a prime ring, then $d_2(r) = 0 \forall r \in R$. That is, $d_2 = 0$.

Corollary (3.1). Let U be a nonzero ideal of R and $a, b \in U$. If $[a, [b, x]]_{\sigma, \tau} = 0, \forall x \in U$, then either $a \in C_{\sigma, \tau}$ or $b \in Z(R)$.

Proof. The map $d_1 : R \rightarrow R$, defined by $d_1(x) = [a, x]$, is a (σ, τ) -derivation and

the map $d_2 : R \rightarrow R$, defined by $d_2(x) = [b, x]$, is a derivation.

Moreover $d_2(U) = [b, U] \subset U$, that is $d_2(U) \subset U$ and

$d_1 d_2(U) = d_2(d_2(U)) = [a, [b, U]]_{\sigma, \tau} = (0)$ by assumption. Hence,

in view of Lemma (3.2), we obtain $d_1 = 0$ or $d_2 = 0$. This implies that

$a \in C_{\sigma, \tau}$ or $b \in Z(R)$.

Theorem (3:1). Let $d_1 : R \rightarrow R$ be a (α, τ) -derivation and $d_2 : R \rightarrow R$ be an (α, α) -derivation and $d_3 : R \rightarrow R$ be an (β, β) -derivation. Such that $d_3\beta = \beta d_3, d_2\beta = \beta d_2, d_1\beta = \beta d_1$ where β is automorphism of R . If $U \neq (0)$ is an ideal of R such that $d_3(U) \subset U$ and $d_1 d_2 d_3(U) = 0$, then either $d_1 = 0$ or $d_2 = 0$ or $d_3 = 0$.

Proof: Let $u, v \in U, uv \in U$. By hypothesis $d_2 d_3(U) = 0$, so

$$\begin{aligned} 0 &= d_1 d_2 d_3(uv) = d_1 d_2 (d_3(u)\beta(v) + \beta(u)d_3(v)) d_1 (d_2(d_3(u)(\beta(v)) + d_2(\beta(u)d_3(v))) = \\ &= d_1 (d_2(d_3(u)\beta(v) + d_1(d_3(u)d_2(\beta(v)) + d_2(\beta(v))\alpha d_3(v) + \alpha\beta(u))d_2(d_3(v))) = \\ &= d_1 (d_2(d_3(u)\alpha\beta(v) + d_1(\alpha(d_3(u))d_2(\beta(v)) + d_1(d_2(\beta(u)\alpha d_3(v)) + d_1(\alpha(\beta(u)d_2(d_3(v))) = \\ &= d_1 (d_2(d_3(u)\sigma(\alpha(\beta(v)) + \tau(d_2(d_3(u)d_1(\alpha(\beta(v)) + d_1(\alpha(d_3(u)\sigma d_2(v) + \tau d_3(u)d_1(d_2(\beta(v) + \\ &+ d_1(\alpha(\beta(u)\sigma d_2(d_3(v)) + \tau(\alpha(\beta(u)d_1(d_2(d_3(v))) = 0 \text{ since } (d_1 d_2 d_3 = 0) \\ &\tau(d_2(d_3(d_3(u)d_1(\alpha(\beta(v)) + \alpha d_1(d_3(d_3(u))\sigma(d_2(\beta(v)) + \tau d_3(d_3(u) + d_1(d_2(\beta(v)) + d_1(d_2(\beta(u)\sigma(\alpha(d_3(v) + \\ &+ \tau d_3(d_3(u) + d_1(d_2(\beta(v)) + d_1(d_2(\beta(u)\sigma(\alpha(d_3(v)) \text{ So,} \end{aligned}$$

$$\sigma(d_2(d_3(d_3(u)d_1(\alpha(\beta(u) + \alpha d_1(d_3(d_3(u))\sigma(d_2(\beta(v))d_3(d_3(u)) = 0$$

$$\tau(d_2(d_3^2(u)d_1(\alpha(\beta(v)) + \alpha d_1(d_3^2(u)\sigma(d_2(\beta(v)))\sigma + \tau d_3^2(u)d_1(d_2(\beta(v))) = 0$$

$$\text{Whereas } \tau d_3^2(u)d_1(d_2(\beta(v))) = 0$$

$$\tau d_3^2(u)d_1(\beta(d_2(v))) = 0, \text{ (using } d_2\beta = \beta d_2)$$

$$\tau d_3^2(u)\beta(d_1(d_2(v))) = 0, \text{ (using } d_1\beta = \beta d_1)$$

since β is isomorphism, hence β^{-1} exists $\tau d_3^2(u)(d_1(d_2(v))) = 0$ By Lemma (2.1) we get $d_3^2(u) = 0 \forall u \in U$ or

$$d_1(d_2(v)) = 0 \forall v \in V \text{ This } d_3^2(U) \neq 0 \text{ or } d_1(d_2(v)) = 0$$

• Suppose $d_3^2(U) \neq 0$, then $d_1(d_2(v)) = 0$ by Lemma(2.2) we get

$$d_1 = 0 \text{ or } d_2 = 0$$

• Suppose if $d_1(d_2(v)) \neq 0$, then $d_3^2(U) = 0$

For any $u, v \in U$, so $uv \in U$, Hence

$$0 = d_3^2(U) = d_3^2(uv) = d_3(d_3(uv)) = d_3(d_3(u)\beta(v) + \beta(u)d_3(v)) =$$

$$= d_3(d_3(u)\beta(\beta(v)) + \beta(d_3(u)d_3(\beta(v)) + d_3(\beta(u))\beta(d_3(v)) + \beta(\beta(v))d_3(d_3(v)))$$

but $d_3^2(U) = (0)$ we get $\beta(d_3(u))d_3(\beta(v)) + d_3(\beta(u))\beta(d_3(v)) = 0$. Also $\beta d_3 = d_3 \beta$, we have $d_3(\beta(u))d_3(\beta(v)) + d_3(\beta(u))d_3(\beta(v)) = 0$, so,

$$2d_3(\beta(u)d_3(\beta(v))) = 0, \text{ since } R \text{ is aprime with ch } R \neq 2 \text{ then}$$

$$d_3(\beta(u)d_3(\beta(v))) = 0, \text{ so, } \beta(d_3(u)d_3(v)) = 0 \text{ and } (d_3(u)d_3(v)) = 0 \forall u, v \in U$$

There fore $d_3(U)d_3(U) = 0$, by Lemma (3.1) we get lither $d_3(U) = (0)$ or $d_3 = 0$

If $d_3(U) = (0)$, then $d_3(ru) = 0, u \in U, r \in R$ this is implies

$$0 = d_3(3)\beta(u) + \beta(r)d_3(u) = d_3(r)\beta(u) \text{ that is}$$

$d_3(r)\beta(u) = 0, \forall u \in U, r \in R$. Now $0 = d_3(v)(ru) = d_3(r)\beta(r)\beta(u) \forall u \in U, r \in R$ (since R is aprime ring) then $d_3(r) = 0 \forall r \in R$ that is $d_3 = 0$

Theorem (3.2) (In general). Let $d_1 : R \rightarrow R$ be (σ, τ) - derivation and $d_i : R \rightarrow R$ be an (α_j, α_j) - derivation such that $d_1\alpha_1 = \alpha_1d_1, d_i\alpha_j = \alpha_jd_i, i = 2, \dots, n, j = n-1, n \geq 2$, Where α_j is an outomorphism of R if $U \neq (0)$ is an ideal of R such that $d_1(U) \subset U$

and $d_1d_2 \dots d_n(U) = (0), n \in N$ Then either $d_1 = 0$ or $d_2 = 0$ or $d_n = 0$

Proof. Let $u, v \in U, uv \in U, d_1d_2 \dots d_n(U) = (0)$, so

$$0 = d_1d_2 \dots d_n(uv) = d_1d_2 \dots d_{n-1}(d_n(u)\alpha_{n-1}(v) +$$

$$+ \alpha_{n-1}(u)\alpha_n(v))d_1(d_2 \dots d_{n-2}(d_{n-1}(u)\alpha_{n-1}(v)) + d_2 \dots d_{n-2}(\alpha_{n-1}(u)d_n(v))) =$$

$$= d_1(d_2(\dots(d_{n-2}(d_{n-1}(u) + \alpha_{n-1}(u)) + d_1(d_{n-1}(u)d_2 \dots d_{n-2}(\alpha_{n-1}(u)) +$$

$$+ d_2(\alpha_{n-1}(v)\alpha_{n-1}d_n(v) + \alpha_{n-1} + \alpha_{n-2}\alpha_{n-1}(u)d_2 \dots d_{n-1}(d_n(v))) =$$

$$= d_1(d_2 \dots d_{n-1}(d_n(u)\alpha_{n-2}\alpha_{n-1}(v) + d_1(\alpha_{n-2}(d_n(u)d_2 \dots d_{n-2}(\alpha_{n-1}(v)) +$$

$$+ d_1(d_2 \dots d_{n-1}(\alpha_{n-1}(u))\alpha_{n-2}d_n(v) + d_1(\alpha_{n-1}(u))d_2 \dots d_{n-1}(d_{n-1}(v))) =$$

$$= d_1(d_2 \dots d_{n-1}(d_n(u)\sigma(\alpha_{n-2}(\alpha_{n-1}(v)) + \tau(d_2 \dots d_{n-1}(d_n(u))d_1(\alpha_{n-2}(\alpha_{n-1}(v)) +$$

$$+ d_1(\alpha_{n-2}(d_{n-1}(u))\sigma(d_2 \dots d_{n-1}(\alpha_{n-1}(v)) + \tau d_n(u)d_1(d_2 \dots d_{n-1}(\alpha_{n-1}(u))\sigma(\alpha_{n-2}(d_n(v)) +$$

$$+ \tau d_2 \dots d_{n-1}(\alpha_{n-1}(u))d_1(\alpha_{n-2}(d_n(u)) + d_1(\alpha_{n-2}(\alpha_{n-1}(u))\sigma d_2 \dots d_{n-2}(d_n(v) +$$

$$+ \tau(\alpha_{n-2}(\alpha_{n-1}(u))d_1(d_2 \dots d_{n-1}d_3(v))) = 0 \text{ since } d_1d_2 \dots d_n = 0$$

$$\tau(d_2 \dots d_{n-1}(d_n(d_n(u)d_1(\alpha_{n-2}(\alpha_{n-1}(v)) + \alpha_{n-2}d_1(d_n(d_n(u))\sigma(d_2 \dots d_{n-1}(\alpha_{n-1}(v)) +$$

$$+ \tau d_n(d_n(u)) + d_1d_2 \dots d_{n-1}(\alpha_{n-1}(v) + d_1(d_2 \dots d_{n-1}(\alpha_{n-1}(u))\sigma(\alpha_{n-2}(d_n(v) +$$

$$+ \tau d_2 \dots d_{n-1}(\alpha_{n-1}(u))d_1(\alpha_{n-2}(d_{n-1}(v) + d_1(\alpha_{n-1}(\alpha_{n-1}(u))\sigma d_2 \dots d_{n-1}(d_n(v))) = 0 \text{ So,}$$

$$\sigma(d_2 \dots d_{n-1}(d_n(d_n(u))d_1(\alpha_{n-2}(\alpha_{n-1}(u)) + \alpha_{n-2}d_1(d_n(d_n(u))\sigma(d_2 \dots d_{n-1}(\alpha_{n-1}(u))d_n(d_n(u))) = 0$$

$$(d_2 \dots d_{n-1}(d_n^2(u)d_1(\alpha_{n-2}(\alpha_{n-1}(u)) + \alpha_{n-2}d_1(d_n^2(u)\sigma(d_2 \dots d_{n-1}\alpha_{n-1}(v)))\sigma + \tau d_{n-1}^2(u)d_1(d_2 \dots d_{n-1}(\alpha_{n-1}(v))) = 0$$

where as

$$\tau d_n^2(u)d_1(d_2 \dots d_{n-1}\alpha_{n-1}(v)) = 0 \text{ (} d_1\alpha_1 = \alpha_1d_1, d_i\alpha_j = \alpha_jd_i, i = 2 \dots n, j = n-1, n \geq 2)$$

$$\tau d_n^2(u)d_1(\alpha_{n-1}(d_2 \dots d_{n-1}(v))) = 0 \text{ (} d_1\alpha_1 = \alpha_1d_1, d_i\alpha_j = \alpha_jd_i, i = 2 \dots n, j = n-1, n \geq 2)$$

$$\tau d_n^2(u)\alpha_{n-1}(d_1(d_2 \dots d_n(v))) = 0 \text{ (} d_1\alpha_1 = \alpha_1d_1, d_i\alpha_j = \alpha_jd_i, i = 2 \dots n, j = n-1, n \geq 2)$$

since α_{n-1}^{-1} is automorphisnhence α_{n-1}^{-1} exist

$\tau d_n^2(u)(d_1(d_2 \dots d_{n-1}(v)) = 0$, by Lemma (3.1) we get $d_n^2(u) = 0$ for all $u \in U$ or $(d_1(d_2 \dots d_{n-1}(v)) = 0$, that is $d_n^2(U) = 0$ or $(d_1(d_2 \dots d_{n-1}(v)) = 0 (n \in N)$

1. Suppose $d_n^2(U) \neq 0$, \Rightarrow then $(d_1(d_2 \dots d_{n-1}(v)) = 0 (n \in N)$

by Lemma (3.2) we get $d_1 = 0$ or $(d_2 \dots d_{n-1}(v)) = 0$

• Suppose $d_1 \neq 0$, \Rightarrow then $(d_2 \dots d_{n-1}(v)) = 0$, by Lemma (2.2)

$d_2 = 0$ or $(d_3 \dots d_{n-1}(v)) = 0$

Thus, is the same way, we get

$d_1 = 0$ or $d_2 = 0 \dots$ or $d_{n-1} = 0$

2. Suppose if $(d_1(d_2 \dots d_{n-1}(v)) \neq 0$, then $d_n^2(U) = 0$ for any $u, v \in U$, so $u \in U$, hence

$$\begin{aligned} 0 &= d_n^2(U) = d_n^2(uv) = d_n(d_n(uv)) = d_n(d_n(u)\alpha_{n-1}(v) + \alpha_{n-1}(u)(d_n(v))) = \\ &= d_n(d_n(u)\alpha_{n-1}(v) + d_n(\alpha_{n-1}(u)d_n(v)) = d_n(d_n(u)\alpha_{n-1}(v) + \alpha_{n-1}(d_n(u)d_n(\alpha_{n-1}(v))) + \\ &+ d_n(\alpha_{n-1}(u)\alpha_{n-1}(d_n(v)) + (\alpha_{n-1}(d_n(u) + d_n(\alpha_{n-1}(v) + \alpha_{n-1}(\alpha_{n-1}(v)))d_n(d_n(v))) \end{aligned}$$

But $d_n^2(U) = 0$ we get $\alpha_{n-1}(d_n(u)d_n(\alpha_{n-1}(v)) + d_n(\alpha_{n-1}(d_n(v))) = 0$

Also $\alpha_{n-1}d_n = d_n\alpha_{n-1}$, we have $d_n(\alpha_{n-1}(u)d_n(\alpha_{n-1}(v)) + d_n(\alpha_{n-1}(u))d_n(\alpha_{n-1}(v)) = 0$,

so $2d_n(\alpha_{n-1}(u)d_n(\alpha_{n-1}(v)) = 0$, since R is aprim with $chR \neq 2$ then

$$d_n(\alpha_{n-1}(u)d_n(\alpha_{n-1}(v)) = 0, \text{ so } \alpha_n(d_n(u)d_n(v)) = 0 \text{ and } (d_n(u)d_n(v)) = 0 \forall u, v \in U$$

There fore $d_n(U)d_n(U) = (0)$ by Lemma(3.1), we get either $d_n(U) = (0)$ or $d_n = 0$

if $d_n(U) = (0)$, then $d_n(ru) = 0, u \in U, r \in R$

This is implies $0 = d_n(r)\alpha_{n-1}(u) + \alpha_{n-1}(r)d_n(u) = d_n(r)\alpha_{n-1}(u)$ that is $d_n(r)\alpha_{n-1}(u) = 0 \forall u \in U, r \in R$, now.

$$0 = d_n(v)\alpha_{n-1}(ru) = d_n(r)\alpha_{n-1}(r)\alpha_{n-1}(u) \forall u \in U, r \in R \text{ (Since } R \text{ is prime ring), then } d_n(r) = 0 \forall r \in R, \text{ that is } d_n = 0$$

Theorem(3.3).Let U be nonzero ideal of R and $a, b, c \in U$, if $[c, [a, [b, x]]]_{\sigma, \tau} = 0, \forall x \in U$, then either

1. $c, a \in C_{\sigma, \tau}$ or $b \in Z(R)$.

2. $c \in C_{\sigma, \tau}$ or $a, b \in Z(R)$.

Proof.The map $d_1 : R \rightarrow R$ define by $d_1(x) = [c, x]_{\sigma, \tau}$ is a (σ, τ) - derivation and the map $d_2 : R \rightarrow R$ defined by

$$d_2(x) = [a, x], \text{ is derivation and the map } d_3 : R \rightarrow R \text{ defined by } d_3(x) = [b, x] \text{ is a derivation moreover}$$

$$d_3(u) = [b, U] \subset U \text{ this is } d_3(u) \subset U \text{ and } d_1 d_2 d_3(u) = d_1(d_2(d_3(u))) = [c, [a, [b, u]]]_{\sigma, \tau} = (0), \text{ by}$$

assumption.

Hence, in view of Lemma (3.2) use obtain

1. If $d_1 = 0$ or $d_2 d_3 = 0$ this is implies that if $d_1 = 0$ this implies $c \in C_{\sigma, \tau}$.

if $d_2 d_3 = 0 \Rightarrow$ by Lemma (2.2) either $d_2 = 0$ or $d_3 = 0$

This implies $a \in C_{\sigma, \tau}$ or $b \in Z(R) \Rightarrow c, a \in C_{\sigma, \tau}$ or $b \in Z(R)$

2. If $d_1 d_2 d_3 = 0$, then $d_1 d_2 = 0$ or $d_3 = 0$ If $d_1 d_2 = 0$ (by Lemma (2.2)) either

$$d_1 = 0 \text{ or } d_2 = 0 \text{ Hence } d_1 = 0 \text{ this implies } c \in C_{\sigma, \tau} \text{ or } a \in Z(a)$$

If $d_3 = 0$ this implies $b \in Z(R)$ So $c \in C_{\sigma, \tau}$ or $a, b \in Z(R)$

Theorem(3.4) (In general).Let U be an nonzero ideal of R and $a_1, a_2, \dots, a_n \in U$, If

$$\left[a_n \left[a_{n-1}, \left[\dots, \left[a_1, x \right] \right] \dots \right] \right]_{\sigma, \tau} = 0 \forall x \in U, \text{ then either}$$

1. $a_n, a_{n-1}, \dots, a_2 \in C_{\sigma, \tau}$ or $a_1 \in Z(R)$.
2. $a_n \in C_{\sigma, \tau}$ or $a_{n-1}, \dots, a_1 \in Z(R)$.

Proof. The map $d_1 : R \rightarrow R$ define by $d_1(x) = [a_n, x]_{\sigma, \tau}$ is a (σ, τ) - derivation and the map $d_2 : R \rightarrow R$ defined by $d_2(x) = [a_{n-1}, x]$, is derivation and the map $d_n : R \rightarrow R$ defined by $d_n(x) = [a_1, x]$ is a derivation moreover $d_n(U) = [a_1, U] \subset U$ this is $d_n(U) \subset U$ and $d_1 d_2 \dots d_n(U) = d_1(d_2(\dots d_n(U))) = [a_n [a_{n-1}, [\dots, [a_1, x]]] \dots]_{\sigma, \tau} = 0 \forall x \in U$ by assumption. Hence, in view of Lemma (3.2) use obtain

1. If $d_1 d_2 \dots d_n = 0$, then $d_1(d_2 \dots d_n) = 0$ this implies that if $d_1 = 0$ this implies $a_n \in C_{\sigma, \tau}$ (1)

or $d_2 \dots d_n = 0$ (by Theorem (3.2)) either $d_2 = 0$ or $d_3 \dots d_n = 0$

if $d_2 = 0$ This implies $a_{n-1} \in C_{\sigma, \tau}$ (2), Similarly $d_{n-1} d_n = 0$ by corollary (3.1)

Then either $a_2 \in C_{\sigma, \tau}$ or $a_1 \in Z(R)$ (3), From (1), (2) and (3) We get $a_n, a_{n-1}, \dots, a_2 \in C_{\sigma, \tau}$ or $a_1 \in Z(R)$.

2. If $d_1 d_2 \dots d_n = 0$, then $(d_1 d_2 \dots d_{n-1}) d_n = 0$ (by Theorem (3.2))

then either $(d_1 d_2 \dots d_{n-1}) = 0$ or $d_n = 0$ implies that if $d_n = 0$ this implies $a_1 \in Z(R)$ (1)

or $(d_1 d_2 \dots d_{n-1}) = 0$, We can write them $(d_1 d_2 \dots d_{n-2}) d_{n-1} = 0$ (by Theorem (3.2)) then either $d_1 d_2 \dots d_{n-2} = 0$ or $d_{n-1} = 0$ implies that if $d_{n-1} = 0$, then this implies $a_2 \in Z(R)$ (2), Similarly $d_1 d_2 = 0$ (by lemma (3.2))

then either $d_1 = 0$ or $d_2 = 0$ by corollary (3.1), then either

$a_n \in C_{\sigma, \tau}$ or $a_{n-1} \in Z(R)$ (3) from (1), (2) and (3) we get $a_n \in C_{\sigma, \tau}$ or $a_{n-1}, \dots, a_1 \in Z(R)$.

References

- [1] Aydin, N. On one sided (σ, τ) - Lie ideals in prime rings. Tr.J. of Math., 21(1997)295-301.
- [2] Aydin, N. Notes on generalized Lie ideals. Analele Universitatii din Timisoara seria, mathematics -informatica-Vol., XXXVI(2), (1999), 7-13.
- [3] Aydin, N. and Kandamar, H. (σ, τ) - Lie ideals in prime rings. Tr.J. of Math, 18(1994), 143-148.
- [4] Aydin, N. and Kaya, K. some generalization in prime ring with (σ, τ) -derivation, Doga-Tr.Math., 3(1992), 1-8.
- [5] Aydin, N. and Kaya, K. some results on generalized Lie ideals, Balikesir University of Math. Symposium Mayıs. (1996)23-26.
- [6] Aydin, N. and Kaya, K. Golbasi, O. some results for generalized Lie ideals with derivation II, Applied Mathematics E-Notes 1(2001), 24-30.
- [7] Aydin, N. and Kaya, K. Golbasi, O. some results on one sided generalized Lie ideals with derivation, Mathematical Notes 2(2002), 83-89.
- [8] Aydin, N. soyturk, M. (σ, τ) - Lie ideals in prime rings with derivation. Tr.J. of Math., 19(1995)239-244.
- [9] Bergen, J, Herstein, I.N. and Kerr, J, W.: Lie ideals and derivation of prime rings. Journal of Algebra 71(1981)259-267.
- [10] Bergen, J, Herstein, I.N. and Lanski, C. Derivation with invertible Values, Can.J.Math. 2(1993), 300-310.
- [11] Bresar, M. Centralizing Mapping and derivation in prime ring, J. of Algebra 156(1993)385-394.
- [12] Burton, David M. Abstract and Linear Algebra, Addison- Wesley Pub. 1972.
- [13] Herstein, I, N, Non commutative rings, Univ. of Chicago The Mathematical Association of America, 1968.
- [14] Herstein, I, N. Topics in ring theory. Univ. of Chicago press, Chicago, 1969.
- [15] Herstein, I, N.: On the Lie structure of an associative ring, Journal Algebra 14(1970)561-571
- [16] Herstein, I, N.: Ring with involution, The University of Chicago press, Chicago 1976.
- [17] Herstein, I.N.: A note on derivation II., 22(4)(1979), 509-511.
- [18] Kaya, K.: On (σ, τ) -derivation of prime ring Doga- Tr.J. of Math., 12(2)(1988)42-45.
- [19] Kaya, K. (σ, τ) -Lie ideals in prime rings, An Univ. Timisoara, Stiinite Mat. 30, No.2-3(1992)251-255.
- [20] Lee, P.H. and Lee, T.K.: Lie ideals in prime ring with derivation. Bull. Inst Math. Acad. Sinica., 11, (1983)75-80.
- [21] Posner, E.C. Derivations in prime rings. Proc. Amer. Math. Soc. 8(1957), 1093-1100.
- [22] Soyurk, K., (σ, τ) -Lie ideals in prime ring with derivation. Doga Tr.J. of Math., 18(1996), 280-283.
- [23] Wallace, S. Prime rings satisfying a generalized polynomial Identity. J. Algebra 12(1996), 576-584.

- [24] Yasuyuki Hirano and Hisao Tominga, some commutativity theorems for prime rings with derivations and deferentially semiprime ring. Math.J.Okayama Univ. 26(1984).101-108.
- [25] K. A. Jassim and Dr. A.A. Hameed, (σ, τ) -Lie Ideals With Derivations In Prime Rings. Department of Mathematics, College of Science, University of Baghdad.Iraq.
- [26] K. A. Jassim, Journal of Al-Nahrain University Vol.15 (4), December, 2012, pp.188-190 Science 18 8 Some Results on (σ, τ) -Left Jordan Ideals in Prime Rings Department of Mathematics, College of Science, University of Baghdad.Iraq

Author Profile

Neamah Majid Kahmmac, I have a bachelor's degree in mathematics, a college of science, the University of Baghdad. Iraq - Baghdad - Jadiriyah complex - near the Ministry of Science and Technology, Currently I am Master student of the Institute of Mathematics and Computer Science of Southern Federal University, 344090, Rostov-on-Don, st. zorge 28, dormitories \mathcal{N}_0 6B, κ. 213, mmathemtic@yahoo.com, 8-938-155-22-96.

Mehdy Hala Shaker, I have a bachelor's degree in computer science, University of Anbar - Computer College - Iraq-Anbar, Currently I am Master student of the Institute of Mathematics and Computer Science of Southern Federal University, 344090, Rostov-on-Don, st. zorge 28, dormitories \mathcal{N}_0 6B, κ. 213, hmprogeram@yahoo.com, 8-938-162-36-94.

