

An in Vitro Study on the Antibacterial Activity of Kefir Milk, Labzyme, Biozyme and Some Antibiotics against *Staphylococcus aureus* and *Escherichia colicauses* Lameness in Broiler Chickens

Abdullah A. Mohammed¹, Auda A. Twaina²

College of Agriculture - University of Basrah. Department of Animal Production

Abstract: Bacterial chondronecrosis with osteomyelitis (BCO) is the most common cause of lameness in commercial broilers. Bacteria entering the blood via translocation from the respiratory system or gastrointestinal tract spread hematogenously to the proximal epiphyseal-physeal cartilage of rapidly growing femora and tibiae, causing BCO. We tested the hypothesis that probiotic kefir milk might attenuate bacterial translocation and thereby reduce the incidence of BCO. This study demonstrates that kefir possesses antibacterial effect against *Staphylococcus aureus* and *Escherichia coli* as compared with Labzyme. Biozyme and some antibiotics, Penicillin, Gentamycin, Tetracycline and Newmycin. The results indicated that the Newmycin was the best significantly ($p < 0.05$) against *Staphylococcus aureus* and *Escherichia coli* bacteria compared with others antibiotics. Labzyme. Biozyme and Kefir. Kefir was a highly significant ($p < 0.05$) to inhibit the growth of *Staphylococcus aureus* and *Escherichia coli* bacteria as compared with Labzyme, Biozyme, Penicillin, and Tetracycline.

Keywords: Kefir fermented milk, probiotic, antibiotics, *Staphylococcus aureus*, *Escherichia coli*

1. Introduction

Worldwide surveys revealed the prevalence of lameness in poultry, especially broiler chickens. In a study conducted in the United Kingdom, the rate of lameness in broiler chickens was 27.6%, of which 3.3% of birds are not fully able to move, and about 12.5 billion chickens suffering from lameness worldwide and similar prevalence rates were estimated in Denmark and the United States. It was observed that 70-90% of these cases were due to Bacterial Chondronecrosis with Osteomyelitis (BCO). (Knowles et al., 2008). BCO is one of the most common causes of lameness in broiler chickens. Symptoms usually begin after the third week of life and may result in increased mortality up to 5%. The disease was first diagnosed in 1972 in Australia and later mentioned in the United States, Canada and Europe. At the end of the 20th century, high rates of BCO were detected in broiler chickens in the United Kingdom, where a group of bacteria causing the disease were isolated, including *Staphylococcus aureus*, *Escherichia coli*, *Salmonella* spp. (Dinev, 2009; Joiner et al., 2005). The negative impact of the lameness on the economics of broiler chickens is clearly observed through the difficulty of movement due to pain, hence the reduction of feed and water consumption, resulting in poor production (Wideman et al., 2012).

The rapid rate of growth and increase of the weights of chicken broilers are among the main causes of BCO. This high rate and rapid growth leads to fractures and fine cracks in the osteocytes. These cracks lead to closure of the blood vessels in the affected area and thus slow flow of blood. (Kense and Landman, 2011). The mucosal disorder of the gastrointestinal tract due to intestinal microflora disturbance and invasion of intestinal pathogens is an additional factor in the occurrence of BCO. Knowles et al., 2008). This

imbalance leads to the transmission of the pathogenic bacteria from the gastrointestinal tract to the bloodstream and the bacteria in the blood vessels settle in the infected area, which in turn leads to blood clotting and therefore the lack of bone supply in the blood leading to necrosis of the bones. In addition, transient pathogens attach to cartilage joints and colonize areas affected by necrosis, making it difficult for immune cells to reach infected areas due to lack of blood flow due to vascular closure. This immune deficiency can be exacerbated by any general immune deficiency which may occur as a result of certain diseases such as Cumboro, environmental factors or the presence of fungal toxins in feed (Wideman et al., 2012). The use of antibiotics may help to reduce the incidence of lameness problems in poultry, but the Europeans and Americans band using antibiotics in poultry fed (Rigobelo et al., 2011; Bitterncourt et al., 2011). Hence, the use of alternative antibiotics as a controlling health problem, including lameness, has been promoted globally. The use of probiotics is an effective alternative that can be used in the prevention of lameness in broiler chickens. The use of probiotic milk kefir is one of the most effective alternatives that can be used to prevent lameness in broiler chicks. (Wideman et al., 2012; Toghyani et al., 2015).

Kefir is a fermented milk drink produced by the actions of bacteria and yeasts contained in kefir grains, and is reported to have a unique taste and unique properties. During fermentation, peptides and exopolysaccharides are formed that have been shown to have bioactive properties. (Farnworth, 2005). The Food and Agriculture Organization (FAO) and the World Health Organization (WHO) have proposed a definition of kefir based on the microbial composition (FAO / WHO, 2001). The microbial kefir contains a variety of "lactic acid bacteria. It is described as a

symbiotic association between lactic and acetic bacteria and yeast and claimed to act against the pathogenic genera *Salmonella*, *Helicobacter*, *Shigella*, *Staphylococcus* and *E. coli* (Arihara, 1990). The function of the microorganisms constituting the kefir may include production of lactic acid, antibiotics and bactericides, which inhibit the growth of undesirable and pathogenic microorganisms. (Angulo, 1992). However scientific interest in kefir is growing due to its health benefits, a limited number of in vitro experimental studies were performed in order to understand the antimicrobial mechanism of kefir's microbial flora (Anar, 2000). It has been reported that kefir causes a bacteriostatic effect against *E. coli* possibly due to competition for nutrients between kefir microbiota and the test strain and/or due to substances that could appear at early stages in the fermentation of milk (Garrote *et al.*, 2000).

Rodrigues, (2005) and Okda *et al.* (2013) assessed the antibacterial activity of kefir against *Salmonella*, *Shigella* and *Staphylococcus* species. Cevikbas *et al.* (1994) studied the antibacterial and antifungal activities of kefir and kefir grain on *Staphylococcus aureus*. The aim of this study was to competitive activity of Kefir, Labzyme, Biozyme, Gentamycin, Pencilin, Tetracyclin and Neomycin in vitro by assessing the reduction of *Staphylococcus aureus* and *Escherichia coli* chicks tibia bone.

2. Material and Methods

Kefir Production: Experiment was conducted in Microbiology Laboratory at the Animal Production Department in the College of Agriculture - University of Basrah, for the period from 10/4/2015 to 7/5/2015 for the purpose of manufacturing a Probiotic of Kefir Milk. was manufactured according to the manufacturing steps mentioned by Auda, (2013) with some modifications to them in six stages, the first phase included the process of fermentation, The microbiology of the kefir grains was developed by using 10 g of these grains in a liter of fresh pasteurized cow's milk in a tight glass with a piece of gauze cloth, and allowed to ferment and incubate for 24 hours and at a temperature (20-25) °C. Then the kefir grains were removed using a plastic filter, to reuse the granules again. Kefir milk was kept in the refrigerator until it was used. Antibacterial activity was tested for the Efficiency of antibacterial agents in the Probiotic of Kefir Milk and compared with the Labzyme, Biozyme, Pencilin, Gentamycin, Tetracyclin, Neomycin against *Staphylococcus aureus* and *Escherichia coli*.

Preparations of bacterial solutions; *Staphylococcus aureus* and *Escherichia coli* were used as test microorganisms which cause Bacterial Chondronecrosis With Osteomyelitis BCO. They were activated in nutrient broth by fermentation at 35°C for 24 hours. A loop full of the bacteria that activated and enriched in nutrient broth were transferred to sterile saline water and emulsified to a turbidity of McFarland 0.5 density. The final bacterial cell concentration approximated to 10⁸/ml with spectrophotometric method.

Testing antimicrobial activity; Antibiotic activity of kefir milk was evaluated using the disk diffusion method as described by Bauer *et al.* (1966). Probiotics Labzyme and

Biozyme and antibiotic Pencilin, Gentamycin, Tetracyclin and Neomycin were used to compare the antimicrobial activity and 10 µg/ml of antibiotics was pipetted on to 5 mm diameter paper disk. 24 and 48 hours fermented kefir were pipette at the amount of 0.1 ml and 1.2 mg/ml as described by Rodrigues *et al.* (2005). The paper disks with antibiotics and experimental kefir were applied to the agar surface previously inoculated with 0.1 ml organism suspension. These plates were incubated at 37°C for 24 hours and the inhibition zones were measured at the end of fermentation period. Experiments were performed in triplicates and mean values were used.

Statistical Analysis; The test for the inhibition of the samples obtained from kefir grains in relation to the Probiotics Labzyme and Biozyme and antibiotic Pencilin, Gentamycin, Tetracyclin and Neomycin was carried out with three repetitions, observing the absorbance readings at 600 nm by spectrophotometry. The averages of the results for each kefir sample in relation to each pathogen were compared statistically using the RLS test at the 5% significance level (SPSS, 2009).

3. Results and Discussion

The table (1) showed significant differences ($p < 0.05$) between the regions of the microbial inhibition zones of the different treatments. The best effect of Neomycin was 19.42, 18.90 mm for the inhibition area. In the *Staphylococcus aureus* and *Escherichia coli* inhibition area respectively. The table also showed no significant differences in the treatment of Kefir milk with Gentamycin in diameter (mm) inhibition of *Staphylococcus aureus*, *Escherichia coli*, while significantly different with Pencilin, Labzyme, Tetracyclin, Biozyme in the diameter of the inhibition of *Staphylococcus aureus* (17.15, 16.5, 16.1 and 12.47 mm respectively) and the diameter of the *Escherichia coli* inhibition area was 14.92, 14.40, 17.37 and 10.05 mm respectively, and the following images 1, 2, 3 and 4 showed diameter (mm) the inhibition area of *Staphylococcus aureus* and *Escherichia coli*.

Several authors have studied the effect of kefir and its microbial flora on the inhibition of microbial activity *in vitro*, in relation to a large variety of Gram-positive and Gram-negative bacteria as well as some fungi. Several studies have demonstrated the antagonistic effect of kefir (Saloff-Coste 1996; Garrote *et al.* 2000; Güven and Göl 2003; Santos *et al.* 2003; Witthuhn *et al.* 2004; Yuksekag *et al.*, 2004) demonstrated that microorganisms isolated from the kefir grains inhibited the growth of *S. aureus*. These authors suggested that organic acids, hydrogen peroxide and other substances were responsible for the inhibition. Pathogenic bacteria such as *Shigella* and *Salmonella* do not grow in the drinks fermented with kefir grains (Koroleva 1988). *Lactobacillus acidophilus* isolated from the kefir showed inhibitory activity toward various Gram (+) and Gram (-) microorganisms (Gilliland and Speck 1977; Apella *et al.* 1992; Gupta *et al.* 1996). Of the different microorganisms isolated from kefir, *Lactococcus* and the acetic acid bacteria were those which presented the maximum inhibitory effect on coliforms (Van, 2001), and they also inhibited the growth of *S. aureus*, *B. cereus*,

Clostridium tyrobutyricum and *L.monocytogenes*. The antagonistic mechanisms of lactic acid bacteria may include the activity of organic acids, hydrogen peroxide (Shahani and Chandan 1979; Juven et al. 1992), diacetyl, bacteriocins and other compounds (Helander et al.1997).The results corroborate the Hypothesis that these chemical compounds presentkefir grains were associated withAntagonistic mechanisms against the pathogens.

In conclusion, Kefir milk can be used hn broiler diets or water as antibacterial

Table 1: Effect of Antimicrobial Activity on the Treatment of Kefir Milk, Labzyme, Biozyme, Antibiotics, Gentamycin, Pencilin, Tetracyclin, Neomycin on Diameter (mm) *Staphylococcus aureus, Escherichia*

Strain	S. aureus	E. coli
Kefir	18.85b±0.26	17.88 b ± 0.38
Labzyme	16.51 d±0.23	14.40 c ± 0.34
Biozyme	12.47 f±0.11	10.05 d± 0.29
Gentamycin	18.70 b ± 0.21	18.025 b ± 0.19
Pencilin	17.15 c ± 0.27	14.925 c ± 0.29
Tetracyclin	16.1 d± 0.39	17.37 b ± 0.53
Neomycin	19.42a± 0.042	18.90 a ± 0.102
Significant level	*	*

(*) Vertically different letters "between the averages of treatments means that there are significant differences (p < 0.05).



Image 1: Diameter (mm) of the inhibition areas *S. aureus*



Image 2: Diameter (mm) of the inhibition areas *E. coli*



Image 3: Diameter (mm) of the inhibition areas *S. aureus*



Image 4: Diameter (mm) of the inhibition areas *E. coli*

References

- [1] Anar S. 2000. Kefir and its properties (kefir veozellikleri).Facul Vet, Uluda. University., 19: 137-140.
- [2] Angulo L., Lopez E., Lema C. 1992. Microflora present in kefir grains of the Galician region (north-west of Spain). Dairy Res., 60: 263-267.
- [3] Apella MC, González SN, Macías MEN, Romero N, Oliver G. 1992. In vitro studies on the inhibition of the growth of Shigellasonnei by *Lactobacillus casei* and *Lactobacillus acidophilus*. J ApplBacteriol.; 73: 480-483.
- [4] Arihara K., Toba T., Adachi, S. 1990. Immunofluorescence microscopic studies on distribution of L. kefirifaciens and L. kefir in kefir grain. Intern J Food Microbiol., 11: 127-134.
- [5] Auda. A.T.2013. Effect of supplementing Different Levels of Kefir Milk to Drinking Water on Productive Performance, Physiological and Immunological Traits of Broiler.MSC.Thesis, Agriculture college. Al Basra University. Iraq.
- [6] Bauer,AW. ; WM. Kirby ; Sherris JC. and Turck M.1966. Antibiotic susceptibility testing by a standardized single disk method.Am J ClinPathol.. 45(4):493- 496.
- [7] Bitterncourt L.C. ; C.C.daSilva ; P. D. S.R. Garcia ; D. C.Z. Donato ; R. de. Albuquerque and Araújo L. F.2011. Influence of a probiotic on broiler performance. R. Bras. Zootec., 40(12) : 2739-2743.
- [8] Cevikbas A., Yemni E., Ezzedenn F. W., Yardimici T. 1994.Antitumoral, antibacterial and antifungal activities of kefir and kefir grain. Phytother Res., 8: 78-82.
- [9] Dinev, I. 2009. Clinical and morphological investigations on the prevalence of lameness associated

- with femoral head necrosis in broilers. *Br. Poult. Sci.* 50:284–290.
- [10] FAO/WHO. 2001. Health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. *Report of a Joint FAO/WHO Expert Consultation on Evaluation of Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria*; FAO/WHO: American Córdoba Park Hotel, Córdoba, Argentina,; pp. 1-34.
- [11] Farnworth E.R. 2005. Kefir – a complex probiotic. *Food Science and Technology Bulletin: Functional Foods* 2: 1–17.
- [12] Garrote G. I., Abraham A. G., De Antoni G. L. 2000. Inhibitory power of kefir: the role of organic acids. *Food Protect.*, 63: 364-369.
- [13] Gilliland SE, Speck ML. 1977. Antagonistic action of *Lactobacillus acidophilus* toward intestinal and foodborne pathogens in associative cultures. *J.FoodProtect.*, 40: 820-823.
- [14] Guven, A. ; A. Guven, and Gulmez, M. 2003. The effect of kefir on the activities of GSH-Px, GST, CAT, GSH and LPO levels in carbon tetrachloride-induced mice tissues. *Journal of Veterinary Medicine B* 50: 412-416.
- [15] Gupta PK, Mital BK, Garg SK. 1996. Inhibitory activity of *Lactobacillus acidophilus* against different pathogens in milk. *J Food Sci Technol.*, 33: 147-149.
- [16] Helander, I.M.; V. A. Wright and Mattila-Sandholm T.M.1997. Potential of lactic acid bacteria and novel antimicrobials against Gram-negative bacteria *Trends in Food Science and Technology*, 8, 146-150.
- [17] Joiner, K. S., F. J. Hoerr, E. van Santen, and S. J. Ewald. 2005. The avian major histocompatibility complex influences bacterial skeletal disease in broiler breeder chickens. *Vet. Pathol.* 42:275–281.
- [18] Juven BJ, Schved F, Linder P. 1992. Antagonistic compounds produced by a chicken intestinal strain of *Lactobacillus acidophilus*. *J of Food Protect.*55: 157-161.
- [19] Kense, M. J. and Landman, W.J.M. 2011. *Enterococcus* infections in broiler breeders and their off spring : molecular epidemiology. *Avian Pathol.* 40, 603–612.
- [20] Knowles, T.G. ; Kestin S.C.; S. N. Haslam ; E. Green ; A. Butterworth ; S.J.Pope ; D. Pfeiffer & Nicol, C.J.2008. Leg disorders in broiler chickens: Prevalence, risk factor and prevention. *Plo S ONE*.3,E 1545.
- [21] Koroleva NS. Technology of kefir and kumys. *BullIntDairy Fed.* 1988; 277: 96-100.
- [22] Okda A. Y. ; H.A. Yehi ; W.I. E Kholly and I.M. Hosny.2013. Antibacterial activity of Kefir cultural metabolites against some food-borne pathogens and their survival in Kefir- Tallaga cheese. *J. Appl. Sci. Res.*, 9(6): 4114-4124.
- [23] Rigobelo C.; P. Maluta and Ávila A. 2011. Performance of broiler chicken fed probiotics supplemented diet. *Arsveterinaria, jaboticabal*, 27(2): 111-115.
- [24] Rodrigues K. L., Caputo L. R. G., Carvalho J. C. T., Evangelista J., Schneedorf, J. M. 2005. Antimicrobial and healing activity of kefir and kefir extract. *Intern. J. Antimicrobial Agents.*, 25: 404-408.
- [25] Saloff-Coste CJ. 1996. Kefir: Nutritional and health benefits of yoghurt and fermented milks. *Danone World Newsletter.*11: 1-7.
- [26] Santos A, san Mauro M, Sanchez A, Torres JM, Marquina D. 2003. The antimicrobial properties of different strains of *Lactobacillus spp.* Isolated from kefir. *SystApplMicrobiol.*, 26: 434-437.
- [27] Shahani, K.M. and R.C. Chandan.1979. Nutritional and healthful aspects of cultured and culture-containing dairy foods. *Journal of Dairy Science*, 62, 1685-1694.
- [28] SPSS. 2009. Statistical Package of Soc. Sci., Ver.18. Appl. Guide. Copy right by SPSS Inc.USA.
- [29] Toghiani A. M. ; S. k. Mosavi ; M. Modaresi and Landy N.2015. Evaluation of kefir as a potential probiotic on growth performance, serum biochemistry and immune responses in broiler chicks. *Animal Nutrition* 1: 305- 309.
- [30] Van WJ. 2001. The inhibitory activity and sensory properties of kefir, targeting the low-income African consumer market. [MSc Thesis] Stellenbosch South Africa: University of Stellenbosch
- [31] Wideman R.F. Jr. ; KR Hamal; J.M. Stark; J. Blankenship; H.Lester ; K.N.Mitchell ;G. Lorenzoni and Pevzner I.2012. Awire –flooring model for including lameness in broiler : Evaluation of probiotic as prophylactic treatment. *poultry science* 91:870-883.
- [32] Witthuhn RC, Schoeman T, Britz T.J. 2004. Isolation and characterization of the microbial population of different South African kefir grains. *Int J Dairy Technol.*; 57: 33-37.
- [33] Yuksekag ZN, Beyatli Y, Aslim B. 2004. Determination of some characteristics coccid forms of lactic acid bacteria isolated from Turkish kefirs with natural probiotic. *LebensmWiss U Technol.*37: 663-667.