

Assessment of Geochemistry of Soils of Iron Ore Belt of Bolani and Barbil Odisha, India: A Factor Model

Dr. B. K. Ratha

Associate Professor in Geology, P. G. Department of Geology, Utkal University, Vani Vihar, Bhubaneswar-751004, Odisha, India
Corresponding Author E-mail: rathabk62[at]gmail.com

Abstract: *Natural processes such as weathering and erosion control the metal load of soils making it suitable for plant growth. However, mining and industrial activity in a region can greatly obliterate this natural balance by accelerated accumulation of undesirable trace metals in soils, rendering it unsuitable for plant growth. Hence, geochemical study of soils of such area is necessary. Factor analysis was carried out on soil geochemical data to establish various processes responsible for metal accumulation. Five factors were retained from post-monsoon soil data. First and second factors are attributed to iron and manganese mining in the area. The third, fourth and fifth factors are attributed to natural processes such as chemical alteration of iron minerals, weathering of rocks and formation of clay minerals.*

Keywords: Geochemistry, Soil, Post-monsoon, Factor analysis, Bolani and Barbil

1. Introduction

Natural processes such as weathering and erosion control the metal load of soils making it suitable for plant growth. However, mining and industrial activity in a region can greatly alter this natural balance by accelerated accumulation of undesirable trace metals in soils, thereby making it unsuitable for plant growth. Alluvial soils are most favorable sinks for heavy metals in a polluted mining environment (Forstner and Wittmann, 1979). Several investigations carried out on soils (Hren *et al.*, 2001; McGregor *et al.*, 1998; Lin and Herbert, 1997; Bullock and Bell, 1997; Clark *et al.*, 2001; and Gabler and Schneider, 2000; Ratha and Nayak, 2005) reveal that the soils and the sediments of the nearby mining area are severely contaminated by metals in most part of the world. Again, several studies (Nair *et al.*, 1990; Bodo, 1989; and Ansari *et al.*, 2000) reveal that rainfall or snowmelt in the upland regions increases the metal load in lower regions. Ansari *et al.* (2000) recorded higher load of metal pollutants in the sediments and soils of Ganga plain after monsoon. Hence, it is necessary to carry out study of metal load in soils and establishes their sources in post-monsoon period. For this purpose soil samples were collected during 2012-14.

Geological Setting

Bolani and Barbil are located in Keonjhar district of Orissa and are known for their iron and manganese deposits. Rocks of the area belong to Iron Ore Group (IOG) comprising mainly Banded Iron Formation (BIF) and associated volcano-sedimentary rocks. The Banded Iron Formation more or less circumscribes the Singhbhum granite batholith. BIF occurrences in Joda-Barbil-Noamundi, Gorumahisani, Daitari are rich in iron ore deposits. These have been classified into 3 major stratigraphic belts such as BIF-I, BIF-II and BIF-III. The last one is exposed in Bonai-Keonjhar belt of Sundergarh-Keonjhar district (Acharya, 1984).

2. Materials and Methods

Soil samples were collected from 27 locations (Fig.1) during post-monsoon periods covering the Bolani and Barbil region. From each location about 4kg sample was collected in polythene packet. Then the samples were dried and sieved. Coarse materials were removed. One gram of <74 μ m size fraction was taken for metal analysis. One gram of soil (<200mesh fraction) was treated with concentrated acid, dissolved in a closed system. About 25ml of aqua-regia (3 parts HNO₃ and 1 part HCl) was used for dissolution. The reduced volume of the sample solution was then treated with HNO₃ and HClO₄ acids in 1:1 proportion and again concentrated to minimum volume. The solution was then extracted with distilled water and the volume was made to 250 ml. The sample solution was then aspirated into atomic absorption spectrophotometer (Perkin Elmer-3100) under optimum operational conditions to measure the concentration of different elements such as Ca, Na, K, Mg, Fe, Mn, Co, Zn, Ni, Cd, Pb, Cr and Cu (Table-1).

3. Results and Discussion

Calcium concentration in soils of Bolani varies from 40 to 900 μ g/g and that of Barbil from 10 to 1420 μ g/g. Magnesium concentration recorded in post-monsoon soil varies from 600 to 5900 μ g/g in Bolani and from 250 to 6750 μ g/g in Barbil. Sodium concentrations recorded in soils range from 140 to 2300 μ g/g in Bolani whereas in Barbil these range from 130 to 4200 μ g/g. Potassium levels in soils of Bolani vary from 267 to 9500 μ g/g whereas in Barbil these range from 1170 to 23500 μ g/g. Fe in soils of both areas varies from 2.12-60.87%. Soils nearer to mines show higher Fe content than samples away from mines. Manganese in soils of Bolani is low and varies from 0.05-0.8% whereas Barbil soils are richer in Mn, which varies from 0.001 to 10.18%. Cu in soils varies from 28 to 80 μ g/g in Bolani and range from 18 to 140 μ g/g in Barbil. Zn in soils varies from 40 to 120 μ g/g in

Bolani and range from 58 to 260 $\mu\text{g/g}$ in Barbil. Ni in soils varies from 140 to 260 $\mu\text{g/g}$ in Bolani and range from 20 to 440 $\mu\text{g/g}$ in Barbil. Co in soils varies from 80 to 180 $\mu\text{g/g}$ in Bolani and range from 13 to 120 $\mu\text{g/g}$ in Barbil. Cr in soils varies from 200 to 1200 $\mu\text{g/g}$ in Bolani and range from 65 to 970 $\mu\text{g/g}$ in Barbil. Pb content in soils is 5 to 40 $\mu\text{g/g}$ in Bolani and range from 7 to 90 $\mu\text{g/g}$ in Barbil. Cd content in soils varies from 140 to 160 $\mu\text{g/g}$ in Bolani and 1 to 7 $\mu\text{g/g}$ in Barbil. Fe, Mn, Cr, Co, Pb and Cd are higher in soil samples than the world averages reported for soils. Cr contamination is more in soils of Bolani whereas Barbil soils are more contaminated by Pb. Mn contamination is found more in soils of Barbil. Soils of Bolani are more contaminated with heavy metals compared to soils of Barbil due to the location of the area nearer to iron ore mining sites.

Factor Analysis

Factor analysis, one of the possible multivariate analyses has been widely used for the interpretation of geochemical data on sediments, soil and water. The technique was first used in geological evaluation by Krumbein (1957). Factor analysis is a useful explanatory tool in multivariate statistical analysis and it can be applied to discover and interpret relations among variables including samples or to test hypotheses. The general purpose of factor analytic technique is to find a way of condensing the information contained in a number of original variables into a smaller set of new composite dimensions with minimum loss of information (Davis, 1986; Ratha and Sahu, 1993; Ratha, 1997). In order to establish the natural processes and the anthropogenic processes responsible for pollution, the R-mode factor analysis with rotation was applied to normalized major and trace element data of soils (Kaiser, 1958; Kumru and Bakac, 2003).

Five factors were retained (Table-2) for interpretation. These five factors explain 82.16% of total variance.

First factor explains 27.14% of variance and is positively loaded on Fe-Ni-Co-Cr-Cd and negatively on Pb. This factor explains the dominant contribution from iron ore mining activity.

Second factor explains 18.73% of total variance and is loaded on Mn-Cu-Zn-Pb suggesting contribution from manganese mining.

Third factor explains 17.02% of total variance and is positively loaded on Ca-Mg suggesting development of clay minerals in the soil. Negative loading on Fe-Mn-Co reveals the formation of goethite containing manganese and cobalt in its matrix.

Fourth factor explains 9.73% of total variance and is negatively loaded on K-Zn. This is a negative unipolar axis suggesting contribution from shaly host rock.

Fifth factor explains 9.54% of total variance and is positively loaded on Na suggesting absorption of sodium in clay matrix and development of montmorillonite in impeded drainage condition.

4. Conclusion

The application of multivariate statistical analysis for the assessment of geochemical data has led to the determination and classification of different sources of metal loads. Analyses of the metals of the soils of the Bolani and Barbil area show that they are highly contaminated due to mining and related industrial activity of the area. Hence, appropriate remedial steps may be taken to prevent contamination of the soils from mining and industrial activity in rainy season.

References

- [1] Acharya, S. (1984). Stratigraphic and structural evolution of the rocks of iron ore basins in Singhbhum-Orissa iron ore province of India. *Ind. Jour. Earth. Sci., CEISM*, pp. 19-28.
- [2] Adriano, D. C. (1986). Trace elements in terrestrial environment: Springer Verlag, 533p.
- [3] Alloway, B. J. (1990). Heavy metals in soils. Blackie, 331p.
- [4] Ansari, A. A., Singh, I. B. and Toboschall, H. J. (2000). Role of monsoon rain on concentration and dispersion patterns of metal pollutants in sediments and soils of Ganga Plain, India. *Environmental Geology*, Vol. 39(3-4), pp. 221-237.
- [5] Bodo, B. A. (1989). Heavy metals in water and suspended particulates from an urban basin impacting Lake Ontario. *The Science of Total Environment*, Vol. 87/88, Elsevier Science Publishers, B.V. Amsterdam, pp. 329-344.
- [6] Bullock, S. E. T. and Bell, F. G. (1997). Some problems associated with past mining at a mine in the Wetbank coal field, South Africa. *Environmental Geology*, Vol. 33(1) pp. 61-71.
- [7] Chakraborty, K. L. and Majumdar, T. (1986). Geological aspects of the BIF of Bihar and Orissa. *Jour. Geol. Soc. India*, Vol. 28, pp. 109-133.
- [8] Chakraborty, K. L. and Majumdar, T. (2002). Some important aspects of the banded iron formation of eastern Indian shield (Jharkhand and Orissa). *Ind. Jour. Geol.*, Vol. 74, Nos. 1-4, pp. 37-47.
- [9] Clark, M. W., Walsh, S. R. and Smith, J. V. (2001). The distribution of heavy metals in an abandoned mining area; a case study of Strauss Pit, the Drake mining area, Australia: Implications for the environmental management of mine sites. *Environmental Geology*, Vol. 40(6), pp. 655-663.
- [10] Davis, J.C. (1986). Statistics and data analysis in geology. John Wiley and Sons, New York, 550p.
- [11] Forstner, U. and Wittmann, G.T.W. (1979). Metal pollution in the aquatic environment. Springer-Verlag, 486p.
- [12] Forstner, U. (1986). Metal speciation in solid wastes - Factors affecting mobility. In: Speciation of metals in water, sediments and soil systems, Landner, L. (Ed.), Workshop proceedings, Sunne, Springer Verlag, pp. 13-41.
- [13] Gabler, H. E. and Schneider, J. (2000). Assessment of heavy metal of flood soils due to mining and mineral processing in the Harz Mountains, Germany. *Environmental Geology*, Vol. 39(7), pp. 774-782.
- [14] Hren, M. T., Chamberlain, C. P. and Magilliganm, F. J. (2001). A combined flood surface and geochemical analysis of metal fluxes in a historical mined region: a case study from the New World Mining District, Montana. *Environmental Geology*, Vol. 40, pp. 1334-1346.
- [15] Jones, H. C. (1934). The iron ores of Bihar and Orissa. *Memoir GSI*, Vol. 63, pp. 167-302.
- [16] Kaiser, H. F. (1958). The varimax criteria for analytic rotation in factor analysis. *Psychometrika*, Vol. 23, pp. 187-200.

[17] Krumbein, W. C. (1957) Comparison of percentage and ratio data in facies mapping. *J. Sediment. Petrol.* Vol.27, 293-297

[18] Kumru, M.N. and Bakac, M. (2003). R-mode factor analysis applied to the distribution of elements in soils from the Aydin basin, Turkey. *Jour. Geochemical Exploration*, Vol. 77, pp. 81-91.

[19] Lin, Z., Herbert (Jr), R. B. (1997). Heavy metal retention in secondary precipitates from a mine rock dump and underlying soil, Dalarna, Sweden. *Environmental Geology*, Vol. 33(1), pp. 1-12.

[20] McGregor, R. G., Blowes, D. W., Jambor, J. L. and Robertson, W. D. (1998). Mobilization and attenuation of heavy metals within a Nickel mine tailings impoundment near Sudbury, Ontario, Canada. *Environmental Geology*, Vol. 36(3-4), pp. 305-319.

[21] Miller, J.R. (1994). The role of fluvial geomorphic processes in the dispersal of heavy metals from mine sites. *Jour. Geochemical Exploration*, Vol. 58, pp. 101-118.

[22] Mohapatra, B. K., Mishra, P. P., Singh, P. P. and Rajeev (2002). Nature and development of lateritoid type manganese ore bodies in the iron ore group of rocks, Bonai-Keonjhar belt, Orissa. *Ind. Jour. Geol.*, Vol. 74, Nos. 1-4, pp. 145-159.

[23] Nair, S. M., Bolchand, A. N. and Nambisan, P. N. K. (1990). Metal concentrations in recently deposited sediments of Cochin Back waters, India. *The Science of Total Environment*, Vol. 97/98, pp. 507-524.

[24] Ratha, B.K and Naik B.K (2005). Pre- and post-monsoon variation of heavy metals in soils of Barbil, Keonjhar district, Orissa and their implications on environment, *Vistas in Geological Research*, Vol. 4, U.U. Spl. Publ., pp 1-10.

[25] Ratha, D.S. and Sahu, B.K. (1993). Seasonal variation of geo-chemical data in sediment samples from two estuaries in Western India. *Indian Jour. Environmental Health*, Vol. 35, pp. 294-300.

[26] Ratha, D.S. (1997). Sediment pollution studies using statistical techniques. *Jour. Indian Association of Sedimentologists*, Vol.16, No.1, pp. 37-61.

[27] Saha, A. K. (1994). Crustal evolution of Singhbhum – North Orissa, eastern India. *Geol. Soc. Ind.*, Mem. 27, 333p

Table 1: Varimax factor matrix for elemental data of post-monsoon soil samples

	(bold values are significant at 95% level of confidence)					
	Factor-1	Factor-2	Factor-3	Factor-4	Factor-5	Communality
Fe	0.721	0.009	-0.557	0.248	-0.107	0.903
Mn	-0.215	0.674	-0.457	0.196	-0.224	0.797
Ca	-0.246	-0.050	0.800	0.182	0.286	0.818
Mg	0.155	-0.176	0.856	-0.026	-0.045	0.791
Na	-0.223	-0.026	0.179	0.028	0.922	0.934
K	-0.327	0.128	-0.006	-0.869	-0.012	0.877
Cu	0.024	0.944	-0.089	-0.107	0.130	0.928
Zn	0.087	0.771	-0.124	-0.497	-0.044	0.867
Ni	0.826	0.134	0.050	-0.022	-0.029	0.704
Co	0.681	0.120	-0.396	0.156	0.073	0.664
Cr	0.785	-0.234	0.048	0.116	-0.227	0.739
Pb	-0.540	0.552	0.247	0.178	-0.348	0.810
Cd	0.813	-0.231	0.206	0.220	-0.216	0.852
Eigen values	3.5288	2.4353	2.2130	1.2649	1.2399	
% of trace	27.1444	18.7328	17.0230	9.7301	9.5376	
Cumulative	27.1444	45.8772	62.9002	72.6303	82.1679	
% of trace						




Figure 1: Location map of soil samples