

Some Log-Type Classes of Estimators using Multiple Auxiliary Information

Shashi Bhushan¹, Raksoni Gupta², Saurabh Singh³, Anoop Kumar⁴

^{1,3,4}Department of Mathematics and Statistics, Dr. Shakuntala Misra National Rehabilitation University, Lucknow, U.P, India

²Department of Statistics, National Post Graduate College, Lucknow, U.P., India

Abstract: In this paper, some classes of log-type estimators using information on two auxiliary variables have been proposed for estimating the population mean of the study variable. It has been shown that these classes of log-type estimators have lesser mean square error under the optimum values of the characterizing scalars as compared to some of the commonly used estimators available in the literature. Further, an extension of the proposed classes using multiple auxiliary information have also initiated in this article. A numerical study is included as an illustration using two auxiliary variables.

Keywords: Multiple auxiliary information, logarithmic function, bias, mean square error

1. Introduction

In sampling theory, it is a popular trend to use auxiliary information to obtain more efficient estimators for the population parameters to increase the precision of the estimator. Estimators obtained using auxiliary information are supposed to be more efficient than the estimators obtained without using auxiliary information. The ratio, regression, product and difference methods take advantage of the auxiliary information at the estimation stage. Many authors like, Pandey and Dubey (1988), Upadhyaya and Singh (1999), Kalidar and Cingi (2003), Singh and Tailor (2003), Singh (2003), Sisodia and Dwivedi (1981), Koyuncu and Kalidar along with many others have proposed various estimators using auxiliary information on various population parameters like coefficient of skewness, kurtosis, variation, standard deviation, correlation coefficient, etc. Sometimes, it is more economical to obtain information on more than one auxiliary information; this would probably help in improving the efficiency of the estimator, used to estimate the parameter under consideration. The literature deals with a wide range of ratio, product, difference and exponential estimators proposed by various renowned authors using multiple auxiliary information (Olkin (1958), Raj (1965), Singh (1967), Shukla (1966), etc.). Recently, Bhushan, Gupta and Pandey (2015) had made the use of logarithmic relationship between the study variable and auxiliary variable, extending the work, in this paper we have made the use of multiple auxiliary variables x 's for estimating the population mean \bar{Y} . The proposed estimators would work in case when the study variable is logarithmically related to the auxiliary variable.

Consider a finite population $U = (U_1, U_2, \dots, U_N)$ of size N from which a sample of size n is drawn according to simple random sampling without replacement (SRSWOR). Let y_i and x_i denotes the values of the study and auxiliary variables for the i th unit, ($i = 1, 2, \dots, N$), of the population. Further let \bar{y} and \bar{x} be the sample means of the study and auxiliary variables, respectively.

We suggest the following new classes of log-type estimators for the population mean \bar{Y} as:

$$T_{r1} = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1}{\bar{X}_1} \right) \right]^{\alpha_1} \left[1 + \log \left(\frac{\bar{x}_2}{\bar{X}_2} \right) \right]^{\alpha_2} \quad (1.1)$$

$$T_{r2} = \bar{y} \left[1 + \beta_1 \log \left(\frac{\bar{x}_1}{\bar{X}_1} \right) \right] \left[1 + \beta_2 \log \left(\frac{\bar{x}_2}{\bar{X}_2} \right) \right] \quad (1.2)$$

$$T_{r3} = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1^*}{\bar{X}_1^*} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2^*}{\bar{X}_2^*} \right) \right]^{\delta_2} \quad (1.3)$$

$$T_{r4} = \bar{y} \left[1 + \theta_1 \log \left(\frac{\bar{x}_1^*}{\bar{X}_1^*} \right) \right] \left[1 + \theta_2 \log \left(\frac{\bar{x}_2^*}{\bar{X}_2^*} \right) \right] \quad (1.4)$$

where,

$$\begin{aligned} \bar{X}_i^* &= a_i \bar{X}_i + b_i \\ \bar{x}_i^* &= a_i \bar{x}_i + b_i \end{aligned} \quad i = 1, 2$$

such that $\alpha_i, \beta_i, \delta_i$ and θ_i are the optimizing scalars, $a_i (\neq 0), b_i$ are either real numbers or functions of the known parameters of the auxiliary variable x_i 's such as the standard deviations S_{x_i} , coefficient of variation C_{x_i} , coefficient of kurtosis $\beta_2(x_i)$ and correlation coefficient $\rho_{x_i x_j}$ of the population. ($i \neq j = 1, 2$)

2. Properties of the Suggested Classes of Log-Type Estimators

In order to obtain the bias and mean square error (MSE), let us consider

$$\bar{y} = \bar{Y}(1 + e_0); \quad \bar{x}_1 = \bar{X}_1(1 + e_1); \quad \bar{x}_2 = \bar{X}_2(1 + e_2)$$

$$\text{with } E(e_0) = E(e_1) = E(e_2) = 0$$

Using the results given in Sukhatme and Sukhatme (1991), we have the following theorems:

Theorem 2.1

The biases of the suggested classes of estimators are given by

$$Bias(T_{r1}) = \lambda_n \bar{Y} \left(\alpha_1 \rho_{yx_1} C_y C_{x_1} + \alpha_2 \rho_{yx_2} C_y C_{x_2} - \alpha_1 C_{x_1}^2 - \alpha_2 C_{x_2}^2 + \alpha_1 \alpha_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} + \frac{\alpha_1^2}{2} C_{x_1}^2 + \frac{\alpha_2^2}{2} C_{x_2}^2 \right) \quad (2.1)$$

$$Bias(T_{r2}) = \lambda_n \bar{Y} \left(\beta_1 \rho_{yx_1} C_y C_{x_1} + \beta_2 \rho_{yx_2} C_y C_{x_2} + \beta_1 \beta_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} - \frac{\beta_1}{2} C_{x_1}^2 - \frac{\beta_2}{2} C_{x_2}^2 \right) \quad (2.2)$$

$$Bias(T_{r3}) = \lambda_n \bar{Y} \left(\delta_1 \nu_1 \rho_{yx_1} C_y C_{x_1} + \delta_2 \nu_2 \rho_{yx_2} C_y C_{x_2} - \delta_1 \nu_1^2 C_{x_1}^2 - \delta_2 \nu_2^2 C_{x_2}^2 + \delta_1 \delta_2 \nu_1 \nu_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} + \frac{\delta_1^2}{2} \nu_1^2 C_{x_1}^2 + \frac{\delta_2^2}{2} \nu_2^2 C_{x_2}^2 \right) \quad (2.3)$$

$$Bias(T_{r4}) = \lambda_n \bar{Y} \left(\theta_1 \nu_1 \rho_{yx_1} C_y C_{x_1} + \theta_2 \nu_2 \rho_{yx_2} C_y C_{x_2} + \theta_1 \theta_2 \nu_1 \nu_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} - \frac{\theta_1}{2} \nu_1^2 C_{x_1}^2 - \frac{\theta_2}{2} \nu_2^2 C_{x_2}^2 \right) \quad (2.4)$$

Theorem 2.2

The mean square errors of the suggested classes of estimators up to the terms of order n^{-1} are given by

$$MSE(T_{r1}) = \lambda_n \bar{Y}^2 \left(C_y^2 + \alpha_1^2 C_{x_1}^2 + \alpha_2^2 C_{x_2}^2 + 2\alpha_1 \rho_{yx_1} C_y C_{x_1} + 2\alpha_2 \rho_{yx_2} C_y C_{x_2} + 2\alpha_1 \alpha_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} \right) \quad (2.5)$$

$$MSE(T_{r2}) = \lambda_n \bar{Y}^2 \left(C_y^2 + \beta_1^2 C_{x_1}^2 + \beta_2^2 C_{x_2}^2 + 2\beta_1 \rho_{yx_1} C_y C_{x_1} + 2\beta_2 \rho_{yx_2} C_y C_{x_2} + 2\beta_1 \beta_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} \right) \quad (2.6)$$

$$MSE(T_{r3}) = \lambda_n \bar{Y}^2 \left(C_y^2 + \delta_1^2 \nu_1^2 C_{x_1}^2 + \delta_2^2 \nu_2^2 C_{x_2}^2 + 2\delta_1 \nu_1 \rho_{yx_1} C_y C_{x_1} + 2\delta_2 \nu_2 \rho_{yx_2} C_y C_{x_2} + 2\delta_1 \delta_2 \nu_1 \nu_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} \right) \quad (2.7)$$

$$MSE(T_{r4}) = \lambda_n \bar{Y}^2 \left(C_y^2 + \theta_1^2 \nu_1^2 C_{x_1}^2 + \theta_2^2 \nu_2^2 C_{x_2}^2 + 2\theta_1 \nu_1 \rho_{yx_1} C_y C_{x_1} + 2\theta_2 \nu_2 \rho_{yx_2} C_y C_{x_2} + 2\theta_1 \theta_2 \nu_1 \nu_2 \rho_{x_1 x_2} C_{x_1} C_{x_2} \right) \quad (2.8)$$

Corollary 2.3

The minimum value of MSE is obtained for the optimum value of α_i given by

$$\alpha_{1(opt)} = \frac{(\rho_{yx_2} \rho_{x_1 x_2} - \rho_{yx_1}) C_y}{(1 - \rho_{x_1 x_2}^2) C_{x_1}} = \beta_{1(opt)} \quad (2.9)$$

$$\alpha_{2(opt)} = \frac{(\rho_{yx_1} \rho_{x_1 x_2} - \rho_{yx_2}) C_y}{(1 - \rho_{x_1 x_2}^2) C_{x_2}} = \beta_{2(opt)} \quad (2.10)$$

$$\delta_{1(opt)} = \frac{(\rho_{yx_2} \rho_{x_1 x_2} - \rho_{yx_1}) C_y}{(1 - \rho_{x_1 x_2}^2) \nu_1 C_{x_1}} = \theta_{1(opt)} \quad (2.11)$$

$$\delta_{2(opt)} = \frac{(\rho_{yx_1} \rho_{x_1 x_2} - \rho_{yx_2}) C_y}{(1 - \rho_{x_1 x_2}^2) \nu_2 C_{x_2}} = \theta_{2(opt)} \quad (2.12)$$

where $\nu_i = a_i \bar{X}_i / (a_i \bar{X}_i + b_i)$

and the minimum value of MSE is

$$MSE_{min}(T_{ri}) = \lambda_n \bar{Y}^2 C_y^2 (1 - R_{y,12}^2) = M \quad (2.13)$$

where $R_{y,12}^2$ is the multiple correlation coefficient between y and x_1, x_2

Note: It is important to note that if the parameters ρ, S_x^2, S_y^2 are not known then they can be estimated by their unbiased and consistent estimators and following Sampath (2005) it can be easily shown that the resultant estimators attain the same MSE given in (2.13).

3. Multivariate extension of the suggested classes of estimators using multiple auxiliary information

Let there are k auxiliary variables then we can use the variables by taking a linear combination of these k estimators of the form given in section 2, calculated for every auxiliary variable separately, for estimating the population mean. Then the estimators for population mean will be defined as,

$$T_{r1}^* = \bar{y} \prod_{i=1}^k \left[1 + \log \left(\frac{\bar{X}_i}{X_i} \right) \right]^{\alpha_i} \quad (3.1)$$

$$T_{r2}^* = \bar{y} \prod_{i=1}^k \left[1 + \beta_i \log \left(\frac{\bar{X}_i}{X_i} \right) \right] \quad (3.2)$$

$$T_{r3}^* = \bar{y} \prod_{i=1}^k \left[1 + \log \left(\frac{\bar{X}_i}{X_i^*} \right) \right]^{\delta_i} \quad (3.3)$$

$$T_{r4}^* = \bar{y} \prod_{i=1}^k \left[1 + \theta_i \log \left(\frac{\bar{X}_i}{X_i^*} \right) \right] \quad (3.4)$$

where $\alpha_i, \beta_i, \delta_i$ and θ_i are the optimizing scalars ($i = 1, 2, \dots, k$)

4. Properties of the adapted classes of estimators using multiple auxiliary information

$$\bar{y} = \bar{Y}(1 + e_0) ; \bar{x}_i = \bar{X}_i(1 + e_i)$$

with $E(e_0) = E(e_i) = 0 ; (i = 1, 2, \dots, k)$

and using the results given in Sukhatme and Sukhatme (1991), we have the following theorems:

For obtaining Biases and Mean square errors(s) of the adapted classes, let us denote

Theorem 4.1

$$Bias(T_{r1}^*) = \lambda_n \bar{Y} \left(\sum_{i=1}^k \alpha_i \rho_{yx_i} C_y C_{x_i} - \sum_{i=1}^k \alpha_i C_{x_i}^2 + \sum_{i \neq j=1}^k \alpha_i \alpha_j \rho_{x_i x_j} C_{x_i} C_{x_j} + \sum_{i=1}^k \frac{\alpha_i^2}{2} C_{x_i}^2 \right) \tag{4.1}$$

$$Bias(T_{r2}^*) = \lambda_n \bar{Y} \left(\sum_{i=1}^k \beta_i \rho_{yx_i} C_y C_{x_i} + \sum_{i \neq j=1}^k \beta_i \beta_j \rho_{x_i x_j} C_{x_i} C_{x_j} - \sum_{i=1}^k \frac{\beta_i}{2} C_{x_i}^2 \right) \tag{4.2}$$

$$Bias(T_{r3}^*) = \lambda_n \bar{Y} \left(\sum_{i=1}^k \delta_i v_i \rho_{yx_i} C_y C_{x_i} - \sum_{i=1}^k \delta_i v_i^2 C_{x_i}^2 + \sum_{i \neq j=1}^k \delta_i \delta_j v_i v_j \rho_{x_i x_j} C_{x_i} C_{x_j} + \sum_{i=1}^k \frac{\delta_i^2}{2} v_i^2 C_{x_i}^2 \right) \tag{4.3}$$

$$Bias(T_{r4}^*) = \lambda_n \bar{Y} \left(\sum_{i=1}^k \theta_i v_i \rho_{yx_i} C_y C_{x_i} + \sum_{i \neq j=1}^k \theta_i \theta_j v_i v_j \rho_{x_i x_j} C_{x_i} C_{x_j} - \sum_{i=1}^k \frac{\theta_i}{2} v_i^2 C_{x_i}^2 \right) \tag{4.4}$$

Theorem 4.2

The mean square errors of the suggested classes of estimators up to the terms of order n^{-1} are given by

$$MSE(T_{r1}^*) = \lambda_n \bar{Y}^2 \left(C_y^2 + \sum_{i=1}^k \alpha_i^2 C_{x_i}^2 + 2 \sum_{i=1}^k \alpha_i \rho_{yx_i} C_y C_{x_i} + 2 \sum_{i \neq j=1}^k \alpha_i \alpha_j \rho_{x_i x_j} C_{x_i} C_{x_j} \right) \tag{4.5}$$

$$MSE(T_{r3}^*) = \lambda_n \bar{Y}^2 \left(C_y^2 + \sum_{i=1}^k \beta_i^2 C_{x_i}^2 + 2 \sum_{i=1}^k \beta_i \rho_{yx_i} C_y C_{x_i} + 2 \sum_{i \neq j=1}^k \beta_i \beta_j \rho_{x_i x_j} C_{x_i} C_{x_j} \right) \tag{4.6}$$

$$MSE(T_{r3}^*) = \lambda_n \bar{Y}^2 \left(C_y^2 + \sum_{i=1}^k \delta_i^2 v_i^2 C_{x_i}^2 + 2 \sum_{i=1}^k \delta_i v_i \rho_{yx_i} C_y C_{x_i} + 2 \sum_{i \neq j=1}^k \delta_i \delta_j v_i v_j \rho_{x_i x_j} C_{x_i} C_{x_j} \right) \tag{4.7}$$

$$MSE(T_{r4}^*) = \lambda_n \bar{Y}^2 \left(C_y^2 + \sum_{i=1}^k \theta_i^2 v_i^2 C_{x_i}^2 + 2 \sum_{i=1}^k \theta_i v_i \rho_{yx_i} C_y C_{x_i} + 2 \sum_{i \neq j=1}^k \theta_i \theta_j v_i v_j \rho_{x_i x_j} C_{x_i} C_{x_j} \right) \tag{4.8}$$

Corollary 4.3

Minimum value of the mean square error is obtained as

$$MSE_{\min}(T_{ri}^*) = \lambda_n \bar{Y}^2 C_y^2 (1 - R_{y,12,\dots,k}^2)$$

where $R_{y,12,\dots,k}^2$ is the multiple correlation coefficient between y and x_1, x_2, \dots, x_k

constants such that $a_i (\neq 0), b_i$ are either real numbers or functions of the known parameters of the auxiliary variable x_i 's such as the standard deviations S_{x_i} , coefficient of variation C_{x_i} , coefficient of kurtosis $\beta_2(x_i)$ and correlation coefficient $\rho_{x_i x_j}$ of the population. ($i \neq j = 1, 2, \dots, k$). Thus

5. Some members of the class of estimators T_{r3} & T_{r4}

a wide range of estimators can be formed using these known population parameters. In this sub-section, some members of the suggested classes of estimators are given for the case of two auxiliary variables only. As an illustration the members belonging to the classes (3.3), (3.4), we describe the members of the class (3.3) only. The members of the remaining (3.4) can be written on similar lines. Some of them are listed below:

It can be easily observed that the classes T_{r3} & T_{r4} are more generalized form of class of estimators in the sense of the

Table 1

Log type estimators T_{r3}	a_1	b_1	a_2	b_2
$T_{r3_1}^* = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1}{\bar{X}_1} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2}{\bar{X}_2} \right) \right]^{\delta_2}$	1	0	1	0
$T_{r3_2}^* = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1 + C_{x_1}}{\bar{X}_1 + C_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2 + C_{x_2}}{\bar{X}_2 + C_{x_2}} \right) \right]^{\delta_2}$	1	C_{x_1}	1	C_{x_2}

$T_{r_{33}}^* = \bar{y} \left[1 + \log \left(\frac{\beta_2(x_1)\bar{x}_1 + C_{x_1}}{\beta_2(x_1)\bar{X}_1 + C_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\beta_2(x_2)\bar{x}_2 + C_{x_2}}{\beta_2(x_2)\bar{X}_2 + C_{x_2}} \right) \right]^{\delta_2}$	$\beta_2(x_1)$	C_{x_1}	$\beta_2(x_2)$	C_{x_2}
$T_{r_{34}}^* = \bar{y} \left[1 + \log \left(\frac{C_{x_1}\bar{x}_1 + \beta_2(x_1)}{C_{x_1}\bar{X}_1 + \beta_2(x_1)} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{C_{x_2}\bar{x}_2 + \beta_2(x_2)}{C_{x_2}\bar{X}_2 + \beta_2(x_2)} \right) \right]^{\delta_2}$	C_x	$\beta_2(x_1)$	C_{x_2}	$\beta_2(x_2)$
$T_{r_{35}}^* = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1 + S_{x_1}}{\bar{X}_1 + S_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2 + S_{x_2}}{\bar{X}_2 + S_{x_2}} \right) \right]^{\delta_2}$	1	S_{x_1}	1	S_{x_2}
$T_{r_{36}}^* = \bar{y} \left[1 + \log \left(\frac{\beta_1(x_1)\bar{x}_1 + S_{x_1}}{\beta_1(x_1)\bar{X}_1 + S_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\beta_1(x_2)\bar{x}_2 + S_{x_2}}{\beta_1(x_2)\bar{X}_2 + S_{x_2}} \right) \right]^{\delta_2}$	$\beta_1(x)$	S_{x_1}	$\beta_1(x_2)$	S_{x_2}
$T_{r_{37}}^* = \bar{y} \left[1 + \log \left(\frac{\beta_2(x_1)\bar{x}_1 + S_{x_1}}{\beta_2(x_1)\bar{X}_1 + S_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\beta_2(x_2)\bar{x}_2 + S_{x_2}}{\beta_2(x_2)\bar{X}_2 + S_{x_2}} \right) \right]^{\delta_2}$	$\beta_2(x_1)$	S_{x_1}	$\beta_2(x_2)$	S_{x_2}
$T_{r_{38}}^* = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1 + \rho}{\bar{X}_1 + \rho} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2 + \rho}{\bar{X}_2 + \rho} \right) \right]^{\delta_2}$	1	ρ	1	ρ
$T_{r_{39}}^* = \bar{y} \left[1 + \log \left(\frac{\bar{x}_1 + \beta_2(x_1)}{\bar{X}_1 + \beta_2(x_1)} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\bar{x}_2 + \beta_2(x_2)}{\bar{X}_2 + \beta_2(x_2)} \right) \right]^{\delta_2}$	1	$\beta_2(x_1)$	1	$\beta_2(x_2)$
$T_{r_{310}}^* = \bar{y} \left[1 + \log \left(\frac{C_{x_1}\bar{x}_1 + \rho}{C_{x_1}\bar{X}_1 + \rho} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{C_{x_2}\bar{x}_2 + \rho}{C_{x_2}\bar{X}_2 + \rho} \right) \right]^{\delta_2}$	C_{x_1}	ρ	C_{x_2}	ρ
$T_{r_{311}}^* = \bar{y} \left[1 + \log \left(\frac{\rho\bar{x}_1 + C_{x_1}}{\rho\bar{X}_1 + C_{x_1}} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\rho\bar{x}_2 + C_{x_2}}{\rho\bar{X}_2 + C_{x_2}} \right) \right]^{\delta_2}$	ρ	C_{x_1}	ρ	C_{x_2}
$T_{r_{312}}^* = \bar{y} \left[1 + \log \left(\frac{\beta_2(x_1)\bar{x}_1 + \rho}{\beta_2(x_1)\bar{X}_1 + \rho} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\beta_2(x_2)\bar{x}_2 + \rho}{\beta_2(x_2)\bar{X}_2 + \rho} \right) \right]^{\delta_2}$	$\beta_2(x_1)$	ρ	$\beta_2(x_2)$	ρ
$T_{r_{313}}^* = \bar{y} \left[1 + \log \left(\frac{\rho\bar{x}_1 + \beta_2(x_1)}{\rho\bar{X}_1 + \beta_2(x_1)} \right) \right]^{\delta_1} \left[1 + \log \left(\frac{\rho\bar{x}_2 + \beta_2(x_2)}{\rho\bar{X}_2 + \beta_2(x_2)} \right) \right]^{\delta_2}$	ρ	$\beta_2(x_1)$	ρ	$\beta_2(x_2)$

6. Comparison with the Available Estimators

In this section a comparison of the suggested classes of estimators with some of the known estimators in terms of biases and mean square error up to order of is given. We consider the following estimators:

(a) Mean per unit estimator

It is an unbiased estimator of population mean and its variance is given by,

$$Var(\bar{y}) = \lambda_n \bar{Y}^2 C_y^2$$

$$MSE(\bar{y}) - M = R_{y,12}^2 C_y^2 \geq 0 \tag{6.1}$$

(b) Ratio estimator $\bar{y}_R = \bar{y} \frac{\bar{X}}{x}$

where x may be chosen as x_1 or x_2

$$MSE(\bar{y}_R) = \lambda_n \bar{Y}^2 (C_y^2 + C_x^2 - 2\rho_{yx} C_y C_x)$$

$$MSE(\bar{y}_R) - M = \lambda_n \bar{Y}^2 [(C_x - \rho_{yx} C_y)^2 + (R_{y,12}^2 - \rho_{yx}^2) C_y^2] \geq 0 \tag{6.2}$$

Since $R_{y,12}^2 \geq \rho_{yx}^2$

(c) Product estimator $\bar{y}_p = \bar{y} \frac{\bar{x}}{X}$

$$MSE(\bar{y}_p) = \lambda \bar{Y}^2 (C_y^2 + C_x^2 + 2\rho_{yx} C_y C_x)$$

$$MSE(\bar{y}_p) - M = \lambda_n \bar{Y}^2 [(C_x + \rho_{yx} C_y)^2 + (R_{y,12}^2 - \rho_{yx}^2) C_y^2] \geq 0 \tag{6.3}$$

Since $R_{y,12}^2 \geq \rho_{yx}^2$

(d) The usual regression estimator $\bar{y}_{lr} = \bar{y} + \beta(\bar{X} - \bar{x})$

$$MSE(\bar{y}_{lr}) = \lambda \bar{Y}^2 C_y^2 (1 - \rho_{yx}^2)$$

$$MSE(\bar{y}_{lr}) - M = \lambda_n \bar{Y}^2 [(R_{y,12}^2 - \rho_{yx}^2) C_y^2] \geq 0 \tag{6.4}$$

Since $R_{y,12}^2 \geq \rho_{yx}^2$

(e) Log-type estimators

$$T_{G1} = \bar{y} \left[1 + \log \left(\frac{\bar{x}}{X} \right) \right]^{\alpha_1}$$

$$T_{G2} = \bar{y} \left[1 + \alpha_2 \log \left(\frac{\bar{x}}{\bar{X}} \right) \right]$$

$$T_{G3} = \bar{y} \left[1 + \log \left(\frac{\bar{x}}{\bar{X}^*} \right) \right]^{\alpha_3}$$

$$T_{G4} = \bar{y} \left[1 + \alpha_4 \log \left(\frac{\bar{x}}{\bar{X}^*} \right) \right]$$

$$MSE_{\min}(T_{Gi}) = \lambda_n \bar{Y}^2 C_y^2 (1 - \rho_{yx}^2) \quad i = 1, 2, 3, 4$$

$$MSE(T_{Gi}) - M = \lambda_n \bar{Y}^2 [(R_{y.12}^2 - \rho_{yx}^2) C_y^2] \geq 0 \quad (6.5)$$

$$\text{Since } R_{y.12}^2 \geq \rho_{yx}^2$$

7. Numerical Illustration

The comparison among these estimators is given in this section using a real data set. The data for this study is taken from [1], District Handbook of Aligarh, India. The population contains 332 villages. A simple random sample 80 villages is taken for the study. We consider the variables Y , X_1 and X_2 as the number of cultivators, area of the village and number of household in the village respectively. We compute the bias and the MSE for all estimators. The following values were obtained using the whole data sets:

$$\bar{Y} = 1093.1, \quad \bar{X}_1 = 181.57, \quad \bar{X}_2 = 143.31$$

$$C_y = 0.7626, \quad C_{x_1} = 0.7684, \quad C_{x_2} = 0.7616$$

$$\rho_{yx_1} = 0.973, \quad \rho_{yx_2} = 0.862, \quad \rho_{x_1x_2} = 0.842$$

Using the above results we have calculated the MSE and PRE for all the estimators in section 2. The PRE for each estimator with respect to the sample mean of a SRS is defined as follows:

$$e(y') = \left[\frac{MSE(\bar{y})}{MSE(y')} \right] * 100$$

Table 2

Estimators	Auxiliary Variables	MSE	Percent Relative Efficiency (PRE)
\bar{y}	None	6593.042	100
\bar{y}_R	x_1	359.1134	1835.922
\bar{y}_R	x_2	1817.305	362.7923
\bar{y}_P	x_1	26214.39	25.15047
\bar{y}_P	x_2	24520.31	26.88809
\bar{y}_{lr}	x_1	351.218	1877.194
\bar{y}_{lr}	x_2	1694.122	389.1717
T_{Gi}	x_1	351.218	1877.194
T_{Gi}	x_2	1694.122	389.1717
T_{ri}	x_1, x_2	309.8479	2127.832

8. Conclusion

The present study extends the idea regarding the effective use of single auxiliary information on variable to the use of multiple auxiliary information. The biases and mean square error of the suggested classes of estimators are obtained up to the terms of order n^{-1} . It has been shown that the suggested classes have minimum MSE(s) as compared to some renowned estimators available in the literature of sampling. The results are quite convincing both theoretically and empirically.

References

- [1] Ahmed, M.S (1995): Some estimation procedures using multivariate auxiliary information in sample surveys, Ph.D, Thesis, Department of Statistics & Operation Research, AMU
- [2] Bhushan, S. et al. (2013): Improved sampling strategies in finite population, Scholar Press, Germany.
- [3] Bhushan, S., Gupta, R. and Pandey, S.K. (2015): Some log-type classes of estimators using auxiliary information (Submitted).
- [4] Crame'r, H.(1945): Mathematical Methods of Statistics, Princeton: N. J.: Princeton University Press.
- [5] Murthy, M. N.(1967): Sampling Theory and Methods, Calcutta: Statistical Publishing Society, 1967.
- [6] Olkin, I.(1958): "Multivariate Ratio Estimation for Finite Populations," Biometrika, 45 pg 154-65.
- [7] Raj, D.(1965): "On a Method of Using Multi auxiliary Information in Sample Surveys," Journal of the American Statistical Association, 60 pg 270-7.
- [8] Rao, P. S. R. S. and Mudholkar, G. S.(1967):"Generalized Multi- variate Estimator for the Mean of Finite Populations," Journal of the American Statistical Association 62 pg 1009-12.
- [9] Shukla, G. K.(1966): "An Alternative Multivariate Ratio Estimate for Finite Population," Calcutta Statistical Association Bulletin 15 pg 127-34.
- [10]Singh, M. P.(1967): "Ratio cum Product Method of Estimation ," Metrika, 12 (No. 1) pg, 34-42.
- [11]Singh, M. P.(1967): "Multivariate Product Method of Estimation for Finite Populations," Journal of the Indian Society of Agri- cultural Statistics, 19 pg 1-10.
- [12]Smith, T. M. F.(1966): "Ratio of Ratios and Their Applications," Journal of the Royal Statistical Society, Ser. A, 129 (Part 4), pg 531-3.
- [13]Srivastava, S. K.(1965): "An Estimate of the Mean of a Finite Population Using Several Auxiliary Variables," Journal of the Indian Statistical Association, 3 pg 189-94.
- [14]Srivastava, S. K.(1967): "An Estimator Using Auxiliary Information in Sample Surveys," Calcutta Statistical Association Bulletin, 16 pg 121-32.
- [15]Bhushan S., Misra P.K. and Yadav S.K. (2017): On the class of double sampling exponential ratio type estimator using auxiliary information on an attribute and an auxiliary variable, International Journal of Computational and Applied Mathematics, 12(1), 1-10.
- [16]Bhushan S., Misra P.K. and Yadav S.K. (2017): A generalized class of unbiased estimators for population mean using auxiliary information on an attribute and an

- auxiliary variable, International Journal of Computational and Applied Mathematics, 12(1), 11-28.
- [17] Bhushan S., Misra P.K. and Yadav S.K. (2017): On the class of double sampling ratio estimators using auxiliary information on an attribute and an auxiliary variable, International Journal of Statistics and Systems, 12(1), 15-23.
- [18] Bhushan S., Misra P.K. and Yadav S.K. (2017): A generalized class of double sampling estimators using auxiliary information on an attribute and an auxiliary variable, International Journal of Computational Intelligence Research, 13(1), 23-33.
- [19] Bhushan S., Misra P.K. and Yadav S.K. (2017): On unbiased class of ratio estimator for population mean using auxiliary information on an attribute and a variable, International Journal of Statistics and Systems, 12(1), 25-32.
- [20] Bhushan S., Misra P.K. and Yadav S.K. (2017): An unbiased class of ratio type estimator for population mean using an attribute and a variable, Advances in Computational Sciences and Technology, 10(1), 139-146.
- [21] Bhushan S., Misra P.K. and Yadav S.K. (2017): An unbiased estimator for population mean using an attribute and an auxiliary variable, Advances in Dynamical Systems and Applications, 12(1), 29-39
- [22] Bhushan S. and Misra P.K. (2017): An improved class of unbiased separate regression type estimator under stratified random sampling, International Journal of Computational Intelligence Research, 13(1), 35-44.
- [23] Bhushan S. and Misra P.K. (2017): On efficient class of estimators for population mean, International Journal of Applied Agricultural Research, 12(1), 13-20.
- [24] Bhushan S. and Misra P.K. (2017): A family of unbiased estimators of population mean using an auxiliary variable, Advances in Computational Sciences and Technology, 10(1), 129-137.
- [25] Bhushan, S. and Gupta, R.(2019): "A Class of Log-Type Estimators for Population Mean Using Auxiliary Information on an Attribute and a Variable Using Double Sampling Technique", International Journal of Computational and Applied Mathematics, Volume 14, Number 1(2019), pp. 1-12
- [26] Bhushan, S. and Gupta, R.(2019): " Searls' Ratio Product Type Estimators ", International Journal of Statistics and Systems, Volume 14, Number 1 (2019), pp. 29-37
- [27] Bhushan, S. and Gupta, R.(2019): " Some Log-Type Classes of Estimators Using Auxiliary Attribute", Advances in Computational Sciences and Technology Volume 12, Number 2 (2019) pp. 99-108
- [28] Bhushan, S. and Gupta, R.(2019): " Some New Log Type Class of Double Sampling Estimators", International Journal of Applied Agricultural Research, Volume 14, Number 1 (2019) pp. 31-40